Museums 2 7991 . 287 v.97

THE ZOOLOGICAL RECORD

VOLUME 97 SECTION 11 1960

THE UNIVERSITY OF MICHIGAN MAR 26 1955

# **TRILOBITA**

COMPILED BY

J. T. TEMPLE, M.A., Ph.D.

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE EIGHT SHILLINGS
1962

It will greatly help in the production of The Zoological Record and assist the Recorders of the individual sections, if authors will forward a copy of their paper or memoir to the Editor of the Zoological Record, The Zoological Society of London, Regent's Park, London, N.W.1. In the case of separately printed copies of articles so forwarded, the original pagination should be given.

All business correspondence concerning the Record should be addressed to the Scientific Director, The Zoological Society of London, Regent's Park, London, N.W.1.

# THE MUSEUMS JOURNAL

Published by the Museums Association 22 Fitzroy Street, Fitzroy Square, London, W.1.

This journal, which is published monthly, contains articles, reviews and technical notes on every Museum aspect of zoology and other sciences. It is the only English journal which deals with the installation and preservation of exhibits, and which reviews Museum work in all parts of the world.

It is published monthly price 4s., by the Association and can be obtained from the above address. It is distributed free to all members of the Museums Association (Subscription £3 3s. 0d. per annum), of which full particulars are obtainable from the Secretary at the above address.

# TRILOBITA

COMPILED BY

J. T. TEMPLE, M.A., Ph.D.

Al schie Mitt. xi–x

Aust bonii logy [Des

An Circ.
An Hum
16 1:
An Spece Lime
18-1
An May
figs.
An An May

Ba Orda Tain 3], Leni pls.

Upp Tain table publ Arks and addi diag deta

# II. TRILOBITA

#### COMPILED BY

# J. T. TEMPLE, M.A., Ph.D.

#### CONTENTS

									PAGE
I.	TITLES	****	****	****	****	****	****	****	3
П.	Subject	INDEX	****	****	****	****	****	****	8
III.	SYSTEMA	TIC INDI	ez.		****	****	****	****	11

# I.—TITLES

Alberti, G. Trilobiten aus den Tentakulitenschiefern (Devon) von Schaderthal (Ost-Thüringen). Mitt. geol. Staatsinst. Hamb. 29 1960: 109-126 pls. xi-xii text-figs. 1-5.

Amos, A. J., Campbell, K. S. W. & Goldring, R. Australosutura gen. nov. (Trilobita) from the Carboniferous of Australia and Argentina. Palaeontology 3 2 1960: 227-236 pls. xxxix-xl text-figs. 1-2 [Description of trilobite by Campbell & Goldring].

Amsden, T. W. (1). Catalog of Hunton Fossils. Circ. Okla. geol. Surv. No. 38 1956: 1-63.

Amsden, T. W. (2). Additional fossils from the Hunton group, Oklahoma. Oklahoma Geology Notes 16 12 1956: 138–139.

Amsden, T. W. & Ham, W. E. A Remarkable Specimen of the Trilobite Isotelus From the Viola Limestone. Oklahoma Geology Notes 19 2 1959: 18-19 fig. 1.

Amsden, T. W. see Sutherland, P. K.

Anderson, J. G. C. The Wenlock Strata of South Mayo. Geol. Mag., Lond. 97 4 1960: 265-275 text-figs. 1-3.

Ankinovich, S. G. see Borukaev, R. A.

Armstrong, A. K. The Mississippian of West-Central New Mexico. Mem. Bur. Min. min. Res. New Mexico 5 1958: 1-32 pls. i-vi text-figs. 1-21.

Balashova, E. A. (1). [Middle and Upper Ordovician and Lower Silurian trilobites of Eastern Taimyr and their stratigraphical significance.] [Part 3]. Sborn. stat. paleont. biostrat. Inst. geol. Arktiki, Leningrad 17 1959 [apparently 1960]: 5-40 [relevant pls. in Ibid. 14] [Russian].

Balashova, E. A. (2). [Trilobites of the Middle and Upper Ordovician and Lower Silurian of Eastern Taimyr.] Leningrad Univ. 1960: 1-110 pls. i-vi tables 1-5 [Essentially the same work as that published in Sborn. stat. paleont. biostrat. Inst. gool. Arktiki 14, 15, 17 1959-1960—recorded Z.R. 96 11 and ebove—with some changes of wording and the addition of pp. 5-19, tables 1-5, and of some specific diagnoses and comparisons p. 57 ff. Not recorded in detail] [Russian].

Balashova, E. A. (3). [A new genus Protoptychopyge from the Lower Ordovician of the Baltic district]. Paleont. Zhurnal, Moscow 1959 4 1960: 125-129 pl. vii (Russian).

Balashova, E. A. see Chernysheva, N. E.

Barlier, J., Le Maître, D. & Rogier, P. Présence des genres Ampyx et Plumulites dans le Gothlandien de la bordure nord-est du Tassili N' Adjer: Adrar Ikohahoène (bassin de Fort de Polignac). C.R. Acad. Sci. Paris 250 26 1960: 4407–4409.

Bassett, D. A. & Walton, E. K. The Hell's Mouth Grits: Cambrian greywackes in St. Tudwal's Peninsula, North Wales. Quart. J. geol. Soc., Lond. 116 1 1960: 85-105 pl. ix text-figs. 1-10 [Trilobita det. Bassett] [Discussion p. 105 by C. J. Stubblefield].

Bel'skaya, T. N. [The late Devonian sea of the Kuznetsk basin, the history of its development, population and sediments]. Trud. paleont. Inst. Akad. Nauk SSR 82 1960: 1-184 pls. i-xvii text-figs. 1-54 tables [Trilobita det. N. E. Chernysheva & Z. A. Maksimova] [Russian].

Bisat, W. S. see Wilson, A. A.

Bogdanov, N. A. & Chugaeva, M. N. [Palaeozoic deposits of the Omulovsky mountains.] Izv. Akad. Nauk SSSR [Geol.] 1960 5: 24–40 figs. 1–6 [Russian].

Bohlin, B. Trilobites of the Genus Megistaspis from the Vaginatum Limestone of Scandinavia. Bull. geol. Instn Univ. Uppsala 38 2 [Publ. palaeont. Instn Univ. Uppsala No. 27] 1960: 155-205 pls. i-xii textfigs. 1-28.

Bordet, P. see Pillet, J.

Borukaev, R. A. et al. [eds.] [Trud. Conference on the unification of the stratigraphical schemes of the Prepalaeozoic and Palaeozoic of Eastern Kazakhstan I]. Akad. Nauk Kazakhstan SSR, Alma-Ata 1960: 1–298 [Contains several articles mentioning trilobites, e.g. Borukaev & Ivshin (q.v.), Chugaeva (q.v.), S. G. Ankinovich: 194–208, I. F. Nikitin: 228–241, T. B. Rukavishnikova: 247–253, K. A. Lisogor: 259–263] [Russian].

Borukaev, R. A. & Ivshin, N. K. (1), [Middle Cambrian of the north-east and east of Central Kazakhstan]. Trud. Conference on the unification of the stratigraphical schemes of the Prepalaeozoic and Palaeozoic of Eastern Kazakhstan I, Akad. Nauk Kazakhstan SSR, Alma-Ata 1960: 117-143 text-figs. 1-2 table [Russian].

Borukaev, R. A. & Ivahin, N. K. (2). [Upper Cambrian and Lower Ordovician of the north-east and east of Central Kazakhstan]. Trud. Conference on the unification of the stratigraphical schemes of the Prepalaeozoic and Palaeozoic of Eastern Kazakhstan I, Åkad. Nauk Kazakhstan SSR, Alma-Ata 1960: 144-174 text-figs. 1-2 tables 1-3 [Russian].

Bouček, B. & Přibyl, A. Revise trilobitů slovenského svrchního karbonu [Revision der Trilobiten aus dem slowakischen Oberkarbon]. Geol. Práce Zprávy, Bratislava 20 1960: 5–16 [Czech], 17–49 [German] pls. i-x.

Boucot, A. J. et al. A late Silurian fauna from the Sutherland River Formation, Devon Island, Canadian Arctic Archipelago. Bull. geol. Surv. Can. 65 1960: i-ix, 1-51 pls. i-x text-figs. 1-10 [Trilobites p. 40 by H. B. Whittington].

Burgess, J. D. see Hoare, R. D.

Bürgi, H. Biostratigrafia de la Sabana de Bogotá y sus alrededores. Bol. geol. Colombia 5 2 1958 : 113-147 pls. i-xix, xx (map).

Campbell, C. J. see McKerrow, W. S.

Campbell, K. S. W. see Amos, A. J.

Carter, E. K. see Noakes, L. C.

Cave, R. A New Species of Tretaspis from South Wales. Geol. Mag., Lond. 97 4 1960: 334-337 pl. x.

Cavet, P. Le Paléozoïque de la zone axiale des Pyrénées orientales françaises entre le Roussillon et l'Andorre (Étude stratigraphique et paléontologique). Bull. Carte géol. Fr. 55 No. 254 1959 : 1-216 [303-518] pls. i-xii text-figs. 1-36 [Trilobita det. J. Pillet].

Cavet, P. see Pillet, J.

Chang Wen-tang. [The Ordovician sediments of China]. Science Record, Peking (ns) 4 4 1960: 217-226 [Russian].

Chernysheva, N. E. et al. Class Trilobita. Trilobites, in Yu. A. Orlov [ed.], Principles of Palaeontology. Arthropods—Trilobitomorphs and Crustaceans [ed. N. E. Chernysheva]. Moscow 1960: 17-194 pls. i-xii text-figs. 1-427 errata slip [Co-authors E. A. Balashova, N. K. Ivshin, N. N. Kramarenko, Z. A. Maksimova, N. V. Pokrovskaya, N. P. Suvorova] [Russian].

Chernysheva, N. E. see Bel'skaya, T. N.

Chlupáč, I. & Vaněk, J. Nové nálezy fauny ve vyšších vrstvách konépruského devonu [Neue Funde der Fauna in den höheren Schichten des Koněpruser Devon]. Čas. Min. Geol., Prague 2 1957 : 349–351 [German summary p. 351] [Photostat only seen].

Chugaeva, M. N. [The stratigraphical significance of the trilobites of the Middle and Upper Ordovician of the Chu-Ili mountains]. Trud. Conference on the unification of the stratigraphical schemes of the Prepalaeozoic and Palaeozoic of Eastern Kazakhstan I, Akad. Nauk Kazakhstan SSR, Alma-Ata 1960: 254-258 text-fig. 1 [Russian].

Chugaeva, N. N. see Bogdanov, N. A.

Churkin, M., Jnr. & Langenheim, R. L., Jnr. Silurian Strata of the Klamath Mountains, California, Amer. J. Sci. 258 4 1960: 258-273 text-figs. 1-4.

Cogney, G. Recherches géologiques au confluent des oueds Bou-Regreg, Grou et Akrech (Marce occidental). Trav. Inst. sci. chérifien [géol. géogr.] § 1957 : 56 pp. pls. i-ii figs. map.

Cowie, J. L. Notes on Lower Cambrian Stratigraphy in the Boreal Regions. Rep. XXI int. geol. Congr. 8 1960: 57-63 text-figs. 1-2.

Crespin, I. Catalogue of Type and Figured Specimens in the Commonwealth Palaeontological Collection, Canberra. Rep. Bur. Min. Res. Aust. No. 54 1960: 1–92 [Trilobita pp. 85–86].

Dean, W. T. (1). The Ordovician trilobite faunas of South Shropshire, I. Bull. Brit. Mus. (nat. Hist.) [Geol.] 4 4 1960: 73-143 pls. xi-xix text-figs. 1-2.

Dean, W. T. (2), The Use of Shelly Faunas in a Comparison of the Caradoc in England, Wales and Parts of Scandinavia. Rep. XXI int. geol. Congr. 7 1960: 82-87.

Dean, W. T. (3). The Ordovician Rocks of the Chatwall District, Shropshire. Geol. Mag., Lond. 97 2 1960: 163-171 text-figs. 1-2.

Dean, W. T. (4). The Silurian Trilobite Dalmanites myops (König). Palaeontology 2 2 1960: 280.

Destombes, J., Hupé, P. & Jeannette, A. L'Ordevicien de la Meseta côtière marcoaine, à l'Est de Casablanea. C.R. Acad. Sci., Paris 249 1959: 1537–1539.

Easton, W. H. Invertebrate Paleontology. Harper & Bros, New York 1960: i-xii, 1-701 text-figs. [Trilobitae pp. 489-519 text-figs. 12.2-12.15].

Ehrenberg, K. Paläozoologie. Springer-Verlag, Vienna 1960 : i-xvi, 1-408 text-figs. 1-175 [Trilobita pp. 92-95 text-figs. 41-43].

Erben, H. K. [Reviews of several articles on trilobites]. Zbl. Geol. [2] 1959 2 1960: 172-197.

Frederickson, E. A. New evidence concerning Dalmanites oklahomae. Oklahoma Geology Notes 20 3 1960: 53-54 figs. 1-2.

Fritz, M. A. Meraspid period (degree 3) of Pseudogygites latimarginatus (Hall). Canad. J. Zool. 37 6 (for 1959) 1960: 1117–1121 pls. i–ii.

Gobbett, D. J. A New Species of Trilobite from the Lower Oslobreen Limestone. Geol. Mag., Lond. 97 6 1960: 457-460 pl. xv text-fig. 6. Oslobr Spitsbe 460 ple

[196

Hen Oslo

203-2

Her Tensle 1960

Hologeol.
maps

Hu

Trilol

25 19

Hu struc Bull. text-

> Hu Ive

regio

Kaz

Marc

Iv Iv Je

of the land text

Pol Soa pew Pal

i-x

Ho 196 i-iv

Ea

an

he

h-

30:

in.

nt

00

ti-

0.

ıd

17

89

n

Cobbett, D. J. & Wilson, C. B. (the late). The Oslobreen Series, Upper Heela Hoek of Ny Friesland, Spitsbergen. Geol. Mag., Lond. 97 6 1960: 441-457, 460 pls. xiii-xiv text-figs. 1-5.

Goldring, R. see Amos, A. J.

Ham, W. E. see Amsden, T. W.

Henningsmoen, G. The Middle Ordovician of the Oslo region, Norway 13. Trilobites of the family Assphidae. Norsk geol. Tidsskr. 40 3-4 1960: 203-258 pls. i-xiv text-figs. 1-5 tables.

Henningsmoen, G. see Stubblefield, C. J.

Hoare, R. D. & Burgess, J. D. Fauna from the Tensleep sandstone in Wyoming. J. Paleont. 34 4 1960: 711-716 pl. xci.

Holtedahl, O. [ed.] Geology of Norway. Norg. geol. Unders. No. 208 1960: 1-540 text-figs. 1-184 maps tables.

Hu, Chung-Hung see Lochman, C.

Hupé, P. (1). Progrès récents dans l'étude des Trilobites. Bull. trimestr. Serv. Inf. Géol., Paris No. 25 1954 : 2-12.

Hupé, P. (2). Réflexions sur quelques traits de la structure céphalique chez les larves de Trilobites. Bull. Soc. géol. Fr. (7) 1 7 (for 1959) 1960: 745-758 text-figs. 1-16.

Hupé, P. R. (3). Sur le Cambrien inférieur du Maroc. Rep. XXI int. geol. Congr. 8 1960 : 75-85 table.

Hupé, P. see Destombes, J.

Ivshin, N. K. [A new biostratigraphical scale of the Upper Cambrian of the Altai-Sayan geosynclinal region.] Trud. Inst. geol. Nauk, Akad. Nauk Kazakhstan SSR, Alma-Ata 3 1960: 3-62 text-figs. 1-2 tables I-III [Russian] [Offprint only seen].

Ivshin, N. K. see Borukaev, R. A.

Ivshin, N. K. see Chernysheva, N. E.

Jeannette, A. see Destombes, J.

Johnson, G. A. L. The Carboniferous Stratigraphy of the Roman Wall district in western Northumberland. Proc. Yorks. geol. Soc. 32 1 1959: 83–130 pl. ii text-figs. 1–11 tables 1–V.

Johnson, G. A. L. see Wilson, A. A.

Kielan, Z. (1). Upper Ordovician trilobites from Poland and some related forms from Bohemia and Scandinavia [Trylobity górno-ordowickie polski i pewne formy zbliżone z Czech i Skandynawii]. Palaeont. polon. No. 11 1959 [1960]: i-vi, 1-198 pls. i-xxxvi text-figs. 1-50 tables.

Kielan, Z. (2). On two Olenellid trilobites from Hornsund, Vestspitsbergen. Studia geol. polon. 4 1960: 83-91 [English], 91-92 [Polish summary] pls. i-iv.

Kobayashi, T. (1). Some Ordovician Fossils from East Tonkin, Viet Nam. Japan. J. Geol. Geogr. 31 1 1960: 39-48 pl. v text-figs. 1-2. Kobayashi, T. (2). The Ordovician of Korea and its Relation to the other Ordovician Territories. Rep. XXI int. geol. Congr. 7 1960: 34-43 text-fig. 1.

Kobayashi, T. (3). The Cambro-Ordovician Formations and Faunas of South Korea, Part VI, Palaeontology V. J. Fac. Sci. Tokyo Univ. [2] 12 2 1960: 217-275 pls. xii-xiv text-figs. 1-10 table.

Kobayashi, T. (4). The Cambro-Ordovician Formations and Faunas of South Korea, Part VII, Palaeontology VI. Supplement to the Cambrian Faunas of the Tsuibon Zone with Notes on some Trilobites Genera and Families. J. Fac. Sci. Tokyo Univ. [2] 12 2 1960: 329-420 pls. xix-xxi text-figs. 1-13.

Kobayashi, T. (5). Studies on Trilobites in Last Three Decades. Trans. palaeont. Soc. Japan (ns) No. 37 1960: 230-237.

Kramarenko, N. N. [A new genus of Monorakeid trilobites from the Ordovician sediments of the Siberian platform.] Doklady Akad. Nauk SSSR (ns) 109 5 1956: 1030-1031 pl. [Russian].

Kramarenko, N. N. see Chernysheva, N. E.

Langenheim, R. L., Jnr. see Churkin, M., Jnr.

Lawson, J. D. The Succession of Shelly Faunas in the British Ludlovian. Rep. XXI int. geol. Congr. 7 1960: 114-125.

Lehmann, U. Salter'sche Einbettung bei Ogygopsis Walc. (Tril.) im Mittel-Kambrium von Britisch-Columbien? Mitt. geol. Staatsinst. Hamb. 29 1960: 104-108 text-figs. 1-2.

Le Maître, D. see Barlier, J.

Linares Rodríguez, A. Nueva especie fósil del Ordoviciense de Ciudad Real. Estud. geol. Inst. Mallada 9 17 1953: 135-136, 137 [French, English, German summaries] pl. xiv.

Lisbôa, M. do A. Uma nova ordem de Trilobites. Rev. Escola de Minas, Ouro Preto 16 4 1951 : 35–38.

Lisogor, K. A. see Borukaev, R. A.

Lochman-Balk, C. Upper Cambrian-Lower Ordovician Boundary in Montana, U.S.A. Rep. XXI int. geol. Congr. 8 1960: 40–46.

Lochman, C. & Hu, Chung-Hung. Upper Cambrian faunas from the northwest Wind River Mountains, Wyoming. Part I. J. Paleont. 34 5 1960: 793-834 pls. xov-0 text-figs. 1-4.

Lu Yen-hao. [The Cambrian sediments of China]. Science Record, Peking (ns) 4 4 1960 : 199-216 [Russian].

McKerrow, W. S. & Campbell, C. J. The stratigraphy and structure of the Lower Palaeozoic rocks of north-west Galway. Sci. Proc. roy. Dublin Soc. [A] 1 3 1960: 27-51 pls. i-vi text-figs. 1-7 tables 1-3.

Maitre, D. Le see Barlier, J.

Majewske, O. P. see Wilson, J. L.

Maksimova, Z. A. [Devonian and Carboniferous trilobites of the Rudny Altai]. Palaeontological basis of the stratigraphy of the Palaeozoic of the Rudny Altai 7 Moscow 1960: 1–122 pls. i–ix text-figs. 1–4 tables [Russian].

Maksimova, Z. A. see Bel'skaya, T. N.

Maksimova, Z. A. see Chernysheva, N. E.

Milošević, V. Prilog za poznavanje trilobitske faune Srbije [Ein Beitrag zur Kenntnis der trilobitischen Fauna von Serbien]. Glasn. Prir. Muz. Beograd [A] 13 1960: 181-184 [Serbo-Croat], 185 [German summary].

Müller, A. H. Lehrbuch der Paläozoologie Band II Invertebraten Teil 2 Mollusca 2-Arthropoda 1. Gustav Fischer, Jena 1960: i-xii, 1-448 text-figs. 1-617 [Klass Trilobita Walch 1771 pp. 350-420 text-figs. 500-615].

Nelson, C. A. Stratigraphic range of Ogygopsis [Abstract]. Bull. geol. Soc. Amer. 71 12 1960: 2070-2071.

Nikitin, I. F. see Borukaev, R. A.

Noakes, L. C., Carter, E. K. & Öpik, A. A. Explanatory Notes on the Urandangi 4-Mile Geological Sheet. 2nd ed. Explan. Notes Bur. Min. Res. Aust. No. 1 1959: 1-16 text-figs. 1-2 table I map. [Trilobita det. Öpik].

Okano, H. see Sugiyama, T.

Okubo, M. [Japanese trilobites.] Earth Science [Chikyū Kagaku], Tokyo No. 4 1951: 133–139 pl. i 2 tables [Japanese and English: offprint (also paginated 25–31) only seen].

Öpik, A. A. & Pritchard, P. W. Cambrian and Ordovician, in D. Hill & A. K. Denmead [eds.], The Geology of Queensland. J. geol. Soc. Australia 7 1960: 89-114 figs. 11-15 map.

Öpik, A. A. see Noakes, L. C.

Orlov, Yu. A. see Chernysheva, N. E.

Orlowski, S. Stratigraphy of Lower Middle Cambrian Beds in the Vicinity of Sandomierz (Central Poland). Rep. XXI int. geol. Congr. 8 1960: 93–101 text-figs. 1–2.

Palmer, A. R. (1). Trilobites of the Upper Cambrian Dunderberg Shale Eureka District, Nevada. Prof. Pap. U.S. geol. Surv. No. 334-C 1960: 53-109 pls. iv-xi text-figs. 5-22.

Palmer, A. R. (2), Stratigraphic range and significance of the Cambrian agnostid genus Glyptagnostus [Abstract]. Bull. geol. Soc. Amer. 71 12 1960: 1941.

Pillet, J. (1). Révision de Acastella rouaulti (de Tromelin et Lebesconte) 1875 (Trilobite, Eodévonien) [Abstract]. C.R. Soc. géol. Fr. 1959 8 1960 : 239.

Pillet, J. (2). Révision de Acastella rouaulti (de Tromelin et Lebesconte) 1875 (Trilobite, Eodévonien). Bull. Soc. géol. Fr. (7) 1 9 (for 1959) 1960: 939-943 pl. xlv text-figs. 1-2. Pillet, J. (3). Étude systématique des Trilobites, in P. Bordet, P. Cavet & J. Pillet, La faune silurienne de Phulohauki près de Kathmandu (Himalaya du Népal). Bull. Soc. géol. Fr. (7) 2 1 1960: 8-12 pl. i text-figs. 2. 4-5.

Pillet, J. see Cavet, P.

Pokrovskaya, N. V. see Chernysheva, N. E.

Poulsen, C. (1). Notes on some Lower Cambrian Fossils from French West Africa. Mat.-fys. Medd. danske Vid. Selsk. 32 7 1960: 1-12 pls. i-iii text-fig. 1.

Poulsen, C. (2). Fossils from the Late Middle Cambrian Bolaspidella Zone of Mendoza, Argentina. Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960: 1-42 pls. i-iii tables.

Prentice, J. E. Dinantian, Namurian and Westphalian rocks of the district south-west of Barnstable, North Devon. Quart. J. geol. Soc., Lond. 115 3 (for 1959) 1960: 261–286 pls. xii–xiii text-figs. 1–7.

Přibyl, A. Nové poznatky o středočeských trilobitech z nadčeledi *Proctacea* (Neue Erkenntnisse übernittelböhmische Trilobiten der Superfamilie *Proctacea*]. Sborn. Úst. úst. geol. **25** [Pal.] (for 1958) 1960: 177–203 [Czech], 204–221 [German] pls. i–iv.

Přibyl, A. see Bouček, B.

Pritchard, P. W. see Öpik, A. A.

Repina, L. N. [Assemblages of trilobites of the Lower and Middle Cambrian of the western part of Eastern Sayan]. Regional Stratigraphy of SSSR, Geol. Inst. Akad. Nauk SSSR 4 1960: 171–232 pls. i-xix tables [Russian].

Rodríguez, A. Linares see Linares Rodríguez, A.

Rogier, P. see Barlier, J.

**Röömusoks**, A. Stratigraphy and Paleogeography of the Ordovician in Estonia. Rep. XXI int. geol. Congr. 7 1960: 58-69 text-figs. 1-2 table 1.

Rozova, A. V. (1). [On some new trilobites from the Upper Cambrian deposits of Salair]. Trud. Inst. geol. geophiz. Akad. Nauk SSSR (Sib. Otdel.), Novosibirsk 1 1960: 201–211 pl. i table [Russian].

Rozova, A. V. (2), [Upper Cambrian trilobites of Salair (Tolstochikhinsky suite)]. Trud. Inst. geol. geophiz. Akad. Nauk SSSR (Sib. Otdel.), Novosibirsk 5 1960: 1-116 pls. i-viii text-figs. 1-10 tables 1-2 [Russian].

Ruggieri, G. Una nuova trilobite del Permiano del Sosio (Sicilia). Mem. Inst. geol. Univ. Padova **21** [6] 1959: 1–9 pl. i text-figs. 1–4.

Rukavishnikova, T. B. see Borukaev, R. A.

Samsonowicz, J. The Lower Cambrian of the Klimontów Anticlinorium. Rep. XXI int. geol. Congr. 8 1960: 86-92 text-fig. 1.

**Sdzuy**, K. (1). Das Kambrium von Deutschland. Rep. XXI int. geol. Congr. **8** 1960 : 102-112 text-fig. 1 table 1. Laun 1960 Bq

Soc.,

[19

brium

the T

tae) | Rozz 223 table

Bt

Sout

Lon

1-8.

pale Serk Srps [Ser

Ber

tex sbin kar Hei

vla

[Cz En 19

for Tr No

DO

8 1

1

[01

6, in

e de

pal).

figs.

rian

edd.

z. 1.

ldle

ina. -42

estble, for

bi-

ber 0e-58)

he

of R,

ls,

A.

m

ŧ.

),

**Sdruy**, K. (2). Zur Wende Präkambrium/Kambrium. Paläont. Z. **34** 2 1960 : 154–160.

Selwood, E. B. Ammonoids and trilobites from the Upper Devonian and lowest Carboniferous of the Launceston area of Cornwall. Palaeontology 3 2 1960: 153-185 pls. xxvi-xxix text-figs. 1-8.

Squirrell, H. C. & Tucker, E. V. The geology of the Woolhope inlier, Herefordshire. Quart. J. geol. Soc., Lond. 116 2 1960: 139-181 pl. xv text-figs. 1-7.

Snajdr, M. Studie o čeledi Scutelluidae (Trilobitae) [A Study of the Family Scutelluidae (Trilobitae)]. Rozpr. Ust. úst. geol. 26 1960: 1–220 [Czech], 223–263 [English summary] pls. i–xxxvi text-figs. 1–61 tables.

Stanton, W. I. The Lower Palaeozoic rocks of South-West Murrisk, Ireland. Quart. J. geol. Soc., Lond. 116 3 1960: 269-291 pls. xviii-xix text-figs. 1-8.

Stevanović, P. M. [Neue in stratigraphischpaleontologischer Hinsicht bedeutende Beiträge aus Serbien, Mazedonien und NO. Bosnien]. Zapis. Srpsk. geol. Drush., Beograd 1951 1954: 45–49 [Serbo-Croat], 49–51 [German summary].

Strnad, V. (1). Odevonských trilobitech z Horního Benešova II [Über Trilobiten im Devon von Horní Benešov II]. Přír. Čas. Slezský 21 3 1960 : 335–349, 354 [Czech], 349–353 [German] pls. i-iv 3 unnumbered text-figs.

Strnad, V. (2). Spodnokarbonští trilobiti ve sbírkách krajského muzea v Olomouci [Die unterkarbonische Trilobiten in den Sammlungen des Heimatkundlichen Museum in Olomouc]. Sborn. vlast. Úst. Olomouc [A] 4 (for 1959) 1960: 101-104 [Czech], 104 [German summary] pl.

Stubblefield, C. J. Trilobites of South West England. Trans. roy. geol. Soc. Cornwall 19 2 (for 1957-8) 1960: 101-112.

Stubblefield, C. J. & Henningsmoen, G. Proposed use of the plenary powers to designate a type-species for the genus *Sphaerocoryphe* Angelin, 1854 (Class Trilobita) in harmony with accepted use. Bull. zool. Nom. 17 6–8 1960: 231–232.

Stubblefield, C. J. see Bassett, D. A.

Sugiyama, T. [On Japanese Trilobites]. Res. Bull. [Stud.] geol. miner. Inst. Tokyo Bunrika Daigaku No. 1 1944: 21–30 pl. i text-figs. table [H. Okano co-author of some taxa] [Japanese].

Sutherland, P. K. & Amsden, T. W. A Re-Illustration of the Trilobite Lonehodomas megeheei Decker From the Bromide Formation (Ordovician) of Southern Oklahoma. Oklahoma Geology Notes 19 10 1959: 212–219 pls. i-ii text-figs. 1-3.

Suvorova, N. P. [Trilobites of the Cambrian of the eastern Siberian Platform. Part 2. Olenellids—Granulariids.] Trud. paleont. Inst. [PIN] SSSR 84 1960: 1-228 pls. i-xvi text-figs. 1-57 tables errata slip [Russian].

Suvorova, N. P. see Chernysheva, N. E.

Tchernysheva, N. E. see Chernysheva, N. E.

Tchugaeva, M. N. see Chugaeva, M. N.

Teixeira, C. L'évolution du territoire portugais pendant les temps anté-mésozoïques. Bol. Soc. geol. Portug. 13 3 1960 : 229-255 pls. i-xvii text-figs. 1-3.

Termier, H. & G. Paléontologie stratigraphique. Masson et Cie, Paris 1960: 1-515 text-figs. 1-3425.

Tillman, C. G. Spathacalymene, an unusual new Silurian trilobite genus. J. Paleont. 34 5 1960: 891–895 pl. exvi.

Tjernvik, T. E. The Lower Didymograptus Shales of the Flagabro Drilling Core. Geol. Fören. Stockh. Förh. 82 2 1960: 203-217 text-figs. 1-3.

Tkachenko, V. B. [ed.] [Stratigraphy of the Sinian and Cambrian sediments of the north-east of the Siberian platform]. Trud. Inst. geol. Arktiki, Leningrad 101 1959: 1-210 text-figs. 1-41 tables 1-8 [Russian].

Tomezykowa, E. Vorläufiger Entwurf der Stratigraphie des oberen und mittleren Ludlows in den Świętokrzyskie Góry. Prager Arbeitstagung über die Stratigraphie des Silurs und des Devons, Üst. úst. geol. Prague 1960: 343–356 text-figs. 1–2 tables 1–2.

Tripp, R. P. Request for a ruling as to whether Lichas aranea Holzapfel, 1895, is to be treated as a junior primary homonym of Lichas araneus Lindström, 1885 (Class Trilobita). Bull. zool. Nom. 17 6-8 1960: 233-234.

Tschernysheva, N. E. see Chernysheva, N. E.

Tschugaeva, M. N. see Chugaeva, M. N.

Tucker, E. V. see Squirrell, H. C.

Vaněk, J. Rozšíření čeledi Calymenidae v středočeském starším paleozoiku (Trilobitae) [On the distribution of the family Calymenidae in the Central Bohemian lower Palaeozoic (Trilobitae)]. Mus. zprávy Pražského kraje 3 5–6 1958: 113–114 pl. x [English summary p. 114] [Photostat only seen].

Vaněk, J. see Chlupáč, I.

Volk, M. Über Trilobiten aus dem Griffelschiefer des Ordoviziums von Steinsch (Thür.) und Umgebung. Geol. Bl. NO-Bayern 10 1 1960: 2–12 text-fig. 1 tables 1–3.

Walton, E. K. see Bassett, D. A.

Whittard, W. F. The Ordovician Trilobites of the Shelve Inlier, West Shropshire Part IV. Palaeontogr. Soc. [Monogr.] 113 (for 1959) 1960: 117-162 pls. xvi-xxi.

Whittington, H. B. (1). Cordania and other trilobites from the Lower and Middle Devonian. J. Paleont. 34 3 1960: 405-420 pls, li-liv text-fig. 1.

Whittington, H. B. (2). Unique Fossils from Virginia. Virginia Minerals 6 3 1960: 1-7 text-figs. 1-5.

W[hittington], H. B. (3). Trilobita. McGraw-Hill Encyclopedia of Science and Technology, New York 14 1960: 102-106 text-figs. 1-4.

Whittington, H. B. see Boucot, A. J.

Wilson, A. A. The Carboniferous Rocks of Coverdale and adjacent valleys in the Yorkshire Pennines. Proc. Yorks. geol. Soc. 32 3 1960: 285–313 pls. vii–ix text-figs. 1–8 [Discussion pp. 315–316 by W. S. Bisat & G. A. L. Johnson].

Wilson, C. B. (the late) see Gobbett, D. J.

Wilson, J. L. & Majewske, O. P. Conjectured Middle Paleozoic History of Central and West Texas. Publ. Univ. Texas Bur. econ. Geol. No. 6017 1960: 65-86 pls. v-vi text-figs. 49-53.

# II.—SUBJECT INDEX

# GENERAL LITERATURE AND HISTORY

General works.—Treatise, Chernysheva et al.; Ontogeny and cephalic segmentation of trilobites, Hupé (2); Scutelluidae, Snajdr; Middle Ordovician Asaphidae of Oslo, Henningsmoen; Trilobites of S.W. England, Stubblefield; Origin of Cambrian faunas, Sdruy (2); Palaeozoic of E. Kazakhstan, Borukaev et al.; Geology of Norway, Holtedahl; Cambrian stratigraphy of China, Lu; Ordovician stratigraphy of China, Chang; Palaeozoic evolution of Portugal, Teixeira; Precambrian & Cambrian of N.E. Siberia, Tkachenko.

Popular works.—Silicified trilobites from Virginia, Whittington (2); Encyclopedia article, Whittington (3).

Textbooks,—Palaeontology, Ehrenberg; Invertebrate palaeontology, Easton, Müller; Stratigraphical palaeontology, H. & G. Termier.

Bibliography.—Relative dates of Barrande 1846 and Beyrich 1846 discussed, Whittard.

Abstracts.—Nelson, Palmer (2), Pillet (1).

Reviews.—Trilobite studies, Hupé (1), Kobayashi (5); Summaries and reviews of several articles, Erben.

Preservation.—Importance of tectonic deformation of fossils stressed, method of obtaining original dimensions outlined, **Henningsmoon**.

Museum Collections.—Hunton fossils (including L. Devonian trilobites) of Oklahoma, Amsden (1), (2); Types etc. in Canberra, Crespin.

Revision of faunas.—Revision of Middle Ordovician Asaphids listed by Bregger 1884, 1887, Holtedahl 1909, Stermer 1953, Henningsmoen; Revision of Middle Cambrian fauna from Cerrillo El Solitario (Argentina) with discussion of generic assignments of Rusconi 1950–52, Poulsen (2). Revision of nomenclature.—Maja Lomovitakaya 1955 renamed Erdelia, Pagetina Lermontova 1940 renamed Neopagetina, Chernysheva et al.

Designation of type species.—Goycoia Rusconi 1950—designation of G. tellecheai R., Poulsen (2).

Selection of neotypes.—Bojoscutellum paliferum (Beyrich), Paralejurus campanifer (B.), Spiniscutellum umbelliferum (B.), Thysanopeltis speciosa (Hawle & Rajdr; Graularia protolenorum Lermontova, Suvorova; Ogygiocaris sarsi Angelin, Henningsmoen; Phyllaspis raniceps Richter 1863, Alberti.

Selection of lectotypes.—Ampyxella edgelli (Reed), Broeggerolithus globiceps (Bancroft), B. harnagensis (B.), B. longiceps (B.), Onnia l'cobboldi (B.), O. gracilis (B.), Smeathenia smeathenensis (B.), Dean (1); Asaphus" nobilis Barrande, Diacanthaspis decacantha (Angelin), Dindymene ornata Linnarsson, Eodindymene pulchra (Olin), Leonaspis centrina (Dalman), Opsimasaphus latus (A.), Pseudosphaerexochus octacanthus (A.), Symphysurus superstes O., Kielan (1); Australosutura gardneri (Mitchell), Amos et al.; Bathyuriscellus robustus Lermontova, Erbia granulosa L., Namanoia namanensis L., Paramicmacca siberica L., Suvorova; Breviscutellum angusticeps (Barrande), B. inerme (B.), Bronteus asperulus H. & C., B. aulicus H. & C., B. berkeleyanus H. & C., B. carinatus H. & C., B. gigas H. & C., B. intermixtus H. & C., B. pulcher H. & C., B. scharyi B., B. sculptus H. & C., B. sparsus H. & C., B. subtriangularis H. & C., Cornuscutellum rhinoceros (B.), Decoroscutellum haidingeri (B.), D. uralicum (Weber), Kolihapeltis brevifrons (B.), K. parabolina (B.), Kosovopeltis partsehi (B.), Meta-scutellum pustulatum (B.), Microscutellum hawlei (B.), M. tenellum (B.), M. zippei (B.), Octobronteus khodalevitchi W., Paralejurus brongniarti (B.), P. dormitzeri (B.), Planiscutellum planum (H. & C.), Platyscutellum spiniferum (B.), P. viator (B.), Poroscutellum porosum spiniferum (B.), F. viator (B.), Foroscuteitum porosum (B.), Protoscutellum simulans (B.), Scabriscutellum caelebs (B.), S. furciferum (H. & C.), S. oblongum (H. & C.), Soutellum edwardsi (B.), "S." schöbli (Novák), "S." sieberi (H. & C.), Snajdr; Cordania becraftensis Clarke, C. cyclurus (Hall & Clarke), Mystrocephala varicella (H. & C.), Whittington (1); Dalmanites vulgaris (Salter), Dean (4); Denemarkia frontalis (Hawle & Corda), Proetus neglectus Barrando, P. signatus Lindström, Přibyl; Dunderbergia nitida (Hall & Whitfield), Pseudagnostus i acutus (Kobayashi), Palmer (1); Griffithides (Bollandia) rozlozeniki Rakusz, Phillipsia collinsi Mitchell, rozlozeniki Rakusz, Phillipeia colisses mucneu, Bouček & Přibyl; Harpes reticulatus Hawle & Corda, Phacops altaicus Tschernyschew, P. cristata pipa Hall & Clarke, P. sternbergi H. & C., Phillipeia ? abscisa Weber, Proetus sibiricus T., Maksimova; Megistaspis acuticauda (Angelin), M. grandis (Sars), M. heros (Dalman), M. lawrowi (Schmidt), Bohlin; Nescuretus ramseyensis Hicks, Ormathops nicholsoni (Salten) Whittard: Onuciocaris dilatata (Brünnich), (Salter), Whittard; Ogygiocaris dilatata (Brünnich), O. d. stromi Angelin, O. sarsi lata Hadding, Hennings-moen; Protoptychopyge prisca (Lamansky), Balashova (3); Spathacalymene nasuta (Ulrich), Tillman.

#### STRUCTURE

Terminology.—Anterior area (bounded laterally by anterior furrows) for area in front of glabella of Nescuretus, Whittard; Paradoublural line for impression of inner edge of doublure on dorsal test, band furrow for furrow delimiting posterior band of occipital and axial thoracic rings, Henningsmoen;

(1); Py pids, Will segment to be sthereby

[1960]

Precrani

thereby glabellas sidered sidered sented l of third

part of found i Henning

> dae, Sn by narr Ante Phillip

Ceph segmen (2); occipit rarely Snajdr

logy,

Cepi pis) h facial mene Corno dorsal surfac

> Fa Brac (1).

Pa cuse Hup Rain

Kiel

glab

each pleu seve form

руg

50

is

Precanidial lobe for part of glabella in front of facial sture in e.g. Encrinurus and Eodindymene, Kielan (1); Pygidial pleural terminology of Brachymetopids, Whittington (1).

Segmentation.—Larval cephalic spines considered to be segmental; cephalic segmentation inferred thereby as follows: ocular, x2, x1, x, pnt, A1, three glabellar lobes, occipital; larval cheek furrows considered intersegmental, Hupé (2); Asaphids considered to show evidence of three segments (x represented by parafrontal band, pnt, antennular) in front of third glabellar lobe, Henningsmoen.

Glabella.—Parafrontal band of Asaphids considered part of glabella; median paired glabellar impressions found in this family may be ligament suspensions, Henningsmoen.

Glabellar furrows.—Alternative interpretation suggested of the furrows (or impressions) of Scutelluide, Snajdr; Furrows of Megistaspis may be occupied by narrow ridges, Bohlin.

Anterior pits.—Two pairs of pits in axial furrows of Phillipsinella parabola, Kielan (1).

Cephalic spines.—Larval cephalic spines considered segmental and a terminology for them suggested, Hupé (2); Some young stages of Scutelluids have an occipital and a median glabellar spine which are only rarely (as in Cornuscutellum) retained in adults, Saaidr.

Anterior cephalic processes.—Discussion of morphology, distribution and function of these processes, description of spatulate process in Spathacalymene, Tillman.

Cephalic sutures. Dindymeninae (except Plasiaspie) have no connective sutures; in Eodindymene facial suture passes in front on to glabella, in Dindymene it lies in marginal furrow, Kielan (1); In Cornovica (Encrinuridae) facial suture is proparian on dorsal surface of test and opisthoparian on ventral surface; connective sutures are present, Whittard.

Ventral cephalic sutures.—Comparison in some U. Cambrian genera, Palmer (1).

Facial sutures.—Present in Cordania and other Brachymetopids except Brachymetopus, Whittington

Alae.—Alae of Megistaspis considered to belong to glabella, Bohlin.

Palpebral lobes.—Bipleurality and tripleurality discussed; meraspis of *Daguinaspis* has undivided lobes, **Hupé** (2).

Rostral plate.—A subtriangular rostral plate occurs in several unrelated genera including Carmon, Kielan (1); Long crescentic rostral plate in Australosutura gardneri (Brachymetopidae), Amos et al.

Pygidium.—Barrande's and Richters' opinion that each pygidial rib in Scutelluidae corresponds to one pleura upheld by discovery of pleural furrows in several spp.; posterior median rib considered to be formed from two fused ribs, Snajdr; In Housia last thoracic segment is often incompletely separated from pygidium, Palmer (1).

Doublure.—Pygidial doublure of Scutelluidae becomes wider during evolution, **Šnajdr**.

Structure of integument.—Canals through test at front of cephalon of Nescuretus, Whittard; Pits and striations of test of Megistaspis, Bohlin.

Ornamentation.—Importance as a specific criterion, Palmer (1); Reticulation in Trinucleids is common in young stages but usually disappears by the adult and its absence cannot be used as a generic criterion, Dean (1).

Musculature.—Muscle insertion areas of Illaenus lacertus, Whittington (3); Externally smooth inner parts of two posterior pairs of glabellar furrows of Platycalymene interpreted as muscle insertions, scars on two posterior pairs of glabellar lobes of Nescuretus similarly interpreted, Whittard.

Teratology.—Structures interpreted as infilled borings in tests of Megistaspis, Bohlin; Pygidium of Dunderbergia bigranulosa with partly fused thoracic segments, pygidium of Housia ovata with right thoracic pleura still fused, asymmetrical cranidium of Minupeltis conservator, Palmer (1); Pygidial deformities in several Scutelluids including healed fractures in Bojoscutellum plaiferum, Breviscutellum inerme and Planiscutellum planum, Snajdr.

### PHYSIOLOGY AND SEX

Dimorphism.—Two types of librigena associated with both Dunderbergia bigranulosa and D. variagranula, Palmer (1); Platycalymene duplicata—Salter's forma femina and forma mas both found preserved in relief, Whittard.

### DEVELOPMENT

Ontogeny.—General work, Hupé (2); Remarks on ontogeny of Ogygiocaris sarsi, Henningsmoen; Immature stages of Kosovopeltis svobadai and other Scutelluids; meraspid stages are characterised by stronger vaulting, narrower glabella, presence of occipital and median glabellar spines; pre-anal hypothesis of origin of segments upheld, Snajdr; Immature stages of Bailiella lantonoisi, Kobayashi (4); Immature stages of Elvinia recemeri, Holcacephalus tenerus (with protaspides), Taenicephalus shumardi, Lochman & Hu; Meraspis (degree 3) of Pseudogygites latimarginatus (Hall), Pritz; Cephalic ontogeny of Onnia superba and Reuscholithus reuschi, Dean (1); Transitory pygidia of Housia ovata, Palmer (1); Young stages of Aldonaia tersa, Bulaiaspis modesta, Elganellus elegans, Malykania nochtujensis, Suvorova.

Ecdysis.—Salter'sche Einbettung described in Ogygopsis but considered in this case not exuvial, Lehmann.

# **EVOLUTION AND CLASSIFICATION**

Evolution.—Neoteny of proparian forms reaffirmed, Hupé (2).

Phylogeny.—Phylogeny of Scutelluidae discussed, Snajdr; Evolution of Shropshire Trinucleidae, Dean (1); Family relations of Redlichioidea and Corynexochoidea, Suvorova.

Evolutionary series.—Series in *Pseudagnostus* showing gradual effacement of posterior part of pygidial axis, Rozova (2).

Postulated lineages.—Suggested lineage in Actinopelties with reduction of genal and pygidial spines, Kielan (1).

Vertical ranges.—Faunas straddling Cambrian-Ordovician junction in Montana, difficulty of drawing this boundary, Lochman-Balk.

Variation.—Variation from 'wide' to 'long' forms of some U. Ordovician spp. considered to be preservational not sexual; no evidence of geographical subspp. in Poland, Bohemia and Sweden, but some (doubtful) examples recognised between Poland and Britain, Kielan (1).

Classification (principles of).—Principles of generic discrimination of Scutelluids discussed, **Snajdr**; Among U. Cambrian trilobites from Nevada species are distinguished primarily on surface ornament, genera on differences of shape, **Palmer** (1); Reticulation not a generic character in Trinucleids, **Dean** (1).

Classification,—Classification into: Miomera (Agnostida) and Polymera (Redlichiida, Corynexochida, Ptychopariida [including Calymenina], Phacopida, Lichida [including Odontopleurina]), Kobayashi (3); Calymenidae and Homalonotidae referred to order Mesoparia, Lishôa.

## ECOLOGY AND HABITS

Habits.—Anterior cephalic processes may have served as counterbalances to keep posterior part of the body slightly raised during feeding, Tillman.

Migration.—Olenellids of Western American affinities in Lower Cambrian (lower Slakli Series) of Spitsbergen, Kielan (2); Migration of Ashgillian faunas between Sweden, Poland, Bohemia and Britain discussed, Kielan (1); At Cerrillo El Solitario (Argentina) there is a mixed Middle Cambrian fauna with Acado-Baltic (Jincella brachymetopa zone) and Pacific (Bolaspidella zone) elements, suggested correlation with the Centropleura vermontensis fauna of Vermont, Poulsen (2); Correlation of U. Cambrian of U.S.A. and U.S.S.R., Jvahin.

# DISTRIBUTION AND FAUNAS

Cambrian.—Australia (Queensland), Öpik & Pritchard; S. Korea, China, Manchuria, Iran, Kobayashi (4); U.S.S.R. (N.E. Siberia), Tkachenko.

Cambrian (Lower).—Boreal regions, Cowie; Britain (N. Wales), Bassett & Walton; French West Africa, Poulsen (1); Morocco (Anti-Atlas), France, Hupé (3); Poland (Holy Cross Mts.), Samsonowicz; Spitsbergen (Vestspitsbergen), Kielan (2).

Cambrian (Lower & Middle).—Germany, Sdzuy (1); U.S.A. (Calif., Nevada), Nelson; U.S.S.R. (E. Sayan), Repina; U.S.S.R. (Siberian Platform), Suvorova.

Cambrian (Middle).—Argentina (Mendoza), Poulsen (2); Australia (Queensland), Noakes et al.; Canada (B.C.), Lehmann; Poland (Holy Cross Mts.), Orlowski; U.S.S.R. (Kazakhstan), Borukaev & Ivshin (1).

Cambrian (Middle & Upper). — U.S.A. (Wyoming), Lochman & Hu.

Cambrian (Upper).—Korea (South), U.S.A., Kobayashi (3); U.S.A., Palmer (2); U.S.A. (Montana), Lochman-Balk; U.S.A. (Nevada), Palmer (1); U.S.S.R. (Altai-Sayan, etc.), Ivshin; U.S.S.R. (Kazakhstan), Borukaev & Ivshin (2); U.S.S.R. (Salair), Rozova (1), (2).

Ordovician.—Eastern Asia, Kobayashi (2); Estonia, Rōōmusoks; Korea (South), Himalayas, China, Sootland, Kobayashi (3); Morocco, Destombes et al.; Spain (Ciudad Real), Linares.

Ordovician (Lower).—Scandinavia, Baltic, Bohlin; Spitsbergen, Gobbett, Gobbett & Wilson; Sweden (Scania), Tjernvik; U.S.A. (Montana), Lochman-Balk; U.S.S.R. (Baltic), Balashova (3); U.S.S.R. (Kazakhstan), Borukaev & Ivshin (2); U.S.S.R. (W. Siberia), Rozova (2).

Ordovician (Lower & Middle).—Australia (Queensland), Öpik & Pritchard; Britain (Shropshire, Wales, Lake District), Whittard.

Ordovician (Middle).—Britain (Shropshire), Dean (1), (3); Britain, Seandinavia, Dean (2); Canada (Quebec), Snajdr; Canada (Ontario), Fritz; Czechoslovakia (Bohemia), Kielan (1); Eire (Mayo), Stanton; Germany (Thuringia), Volk; Norway (Oslo), Swedea, Henningsmoen; U.S.A. (Oklaboma), Amsden & Ham, Sutherland & Amsden; U.S.A. (Virginia), Whittington (2); U.S.S.R. (E. Siberia), Bogdanov & Chugaeva.

Ordovician (Middle or Upper).—Viet Nam, Kobayashi (1).

Ordovician (Middle & Upper),—U.S.S.R. (Arctic), Balashova (1); U.S.S.R. (Kazakhstan), Chugaeva,

Ordovician (Upper).—Britain (S. Wales, N. England), Cave; France (Pyronees), Cavet; Poland (Lysa Gora), Scandinavia, Bohemia, Britain, Kielan (1); U.S.S.R. (Siberia), Kramarenko.

Silurian.—Britain (W. England), Squirrell & Tucker; Czechoslovakia, U.S.S.R., Snajdr; France (Pyrenees), Cavet; U.S.A. (Texas), Wilson & Majewake.

Silurian P.—Nepal, Pillet (3).

Silurian (Lower).—Britain (Eire), McKerrow & Campbell; Sahara, Barlier et al.; U.S.S.R. (Arotio), Balashova (1).

Silurian (Middle).—Eire (Mayo), Anderson; U.S.A. (Indiana), Tillman.

Silurian (Middle or Upper).—U.S.A. (Calif.), Churkin & Langenheim.

Silurian (Middle & Upper).—Britain, Dean (4); Czechoslovakia (Bohemia), Přibyl.

Silurian (Upper).—Britain, Lawson; Canada (Arctic), Boucot et al.; Poland (Lysa Gora), Tomezy-kowa.

Devonian.—Britain (S.W. England), Stubblefield; U.S.A. (Texas), Wilson & Majewske; U.S.S.R. (Altai), Maksimova. Boher Whitti

[1966

Pillet (

slovaki Strnad (Thüri

Deve wood;

Dev

Japan Cari (N.S.V Burge

Car Wilso Czech Arms

> Car Přiby Steva Per

Car

berla

N.
the †
fossil

Ac Leon

Deve Deve Alta

A mac

A pl. i 1880

A. Ma ]

R.

un), 078

nia,

al.:

in; len

R.

ns-

da

n;

g-

78.

ın-

e),

an

),

);

Devonian (Lower).—France (Armorican massif), Pillet (2); U.S.A. (Oklahoma), Frederickson.

Devonian (Lower & Middle).—Czechoslovakia (Bohemia), Přibyl, Šnajdr; Morocco, Cogney; U.S.A., Whittington (1).

Devonian (Middle).—Colombia, Bürgl; Czechoalovakia, Chlupáč & Vaněk; Czechoslovakia (Silesia), Strnad (1); France (Pyrenees), Cavet; Germany (Thüringia), Alberti; Japan (Iwate), Okubo.

Devonian (Middle & Upper).-Japan, Sugiyama.

Devonian (Upper),—Britain (W. England), Selwood; U.S.S.R. (Kuznetsk basin), Bel'skaya.

Devonian (Upper) or Carboniferous (Lower).—Japan (Iwate), Okubo.

Carboniferous.—Argentina (Chubut), Australia (N.S.W.), Amos et al.; U.S.A. (Wyoming), Hoare & Bursess.

Carboniferous (Lower).—Britain (N. England), Wilson; Britain (W. England), Prentice, Selwood; Czechoslovakia, Strnad (2); U.S.A. (N. Mex.), Armstrong; U.S.B.R. (Altai), Maksimova.

Carboniferous (Lower-Upper).—Britain (Northumberland), Johnson.

Carboniferous (Upper).—Czechoslovakia, Bouček & Přibyl; Jugoslavia, Milošević; Jugoslavia (W. Serbia), Stevanović.

Permian (Middle).-Sicily, Ruggieri.

#### III. SYSTEMATIC INDEX

N.B.—Since all the species of the group are fossil, the † used elsewhere in the Zoological Record to indicate fossils is here omitted.

Acadoparadoxides marocanus n. sp. [nom. nud.], A. sacheri (Barr.) p. 81, Hupé (3).

Acanthaloma mirka Marek 1952 a synonym of Leonaspis olini Troedsson p. 101, Kielan (1).

Acanthaloma (A.) tenella sp. n. p. 68 pl. viii fig. 14 Devonian (Upper) U.S.S.R. (Altai), Maksimova Devonian and Carboniferous trilobites of the Rudny Altai 1960.

Acanthalomina see Leonaspis.

Acanthomicmacca Hupé a synonym of Paramicmacca Lermontova p. 68, Suvorova.

Acanthoparypha text-figs. 5A-I, J (reconstr.), Whittington (2).

Acanthopyge aff. haueri (Barrande 1852) p. 64 pl. iii fig. 14 text-fig. 3, A. aff. contusa (Hall & Clarke 1888) p. 66 pl. iii fig. 6, A. ? sp. p. 67 pl. vi fig. 10; A. parva (Khalfin 1948) [not seen] discussed p. 65, Maksimova.

Acanthopyge aff. haueri (Barr.) p. 350, Chlupáč & Vaněk.

Acaste dayiana R. & E. Richter p. 350, Tomczy-kowa.

Acastella macrocentrus Reed, A. spinosa (Salter) p. 177, Squirrell & Tucker.

Acastella rouaulti—revision in abstract p. 239, Pillet (1).

Acastella rouaulti (de Tromelin et Lebesconte) 1875 em. Péneau 1928 description and affinities p. 939 pl. xlv figs. 1-7 text-figs. 1 (reconstr.)-2, Pillet (2).

Acastella spinosa (Salter) p. 350 [altered by hand to A. tiro in offprint from author p. 352], A. sp. p. 350, Tomczykowa,

Achatella see Pterygometopus.

Acidaspis myops Richter 1863 type of Leonaspis (Grossia) subgen. n. p. 116, Alberti.

Acrocephalaspis [Ivshin] may belong to Housiidae p. 74, Palmer (1).

Acrocephalitidae see Elatelimbus gen. n.

Acrocephalitinae discussed with list of genera and referred to Dokimocephalidae p. 383, Kobayashi (4).

Acrolichas clarus Koroleva [not seen] pl. ix fig. 10, Chernysheva et al.

Actinopeltis Hawle & Corda 1847 discussed pp. 49-50, diagrammatic comparison of spp. forming a possible lineage text-fig. 8, A. barrandei sp. n. for Cheirurus insocialis of Barrande 1872 [non 1852] p. 130 text-fig. 8d Bohemia, A. cf. gryphus (B. 1872) p. 131 pl. xxiv fig. 3, A. g. ventral sutures text-fig. 7, A. cf. globosa (B. 1852) p. 131 pl. xxiv figs. 1-2, A. sp. a p. 132 pl. xxiv fig. 4 text-fig. 36, A. sp. b p. 133 pl. xxiii fig. 7 Poland (Lysa Gora) Ordovician (Upper); Cheirurus neglectus of B. 1872 partim [pl. 3 fig. 18] probably belongs to A. insocialis p. 140, Kielan Palaeont. polon. No. 11 1960.

Aeglina ? oblongula Angelin type refigured and a synonym of Microparia speciosa [q.v.] p. 89, Kielan (1).

Aglagalia n. gen. [nom. nud.] [Termierellidae] p. 77, A. brevissima n. sp. [nom. nud.] p. 81, Hupé (3).

Agnostid indt. p. 40 pl. v fig. 3, Kobayashi (1).

 $\begin{array}{ll} {\bf Agnostidae \ see \ Aspidagnostinae \ and \ Clavagnostinae \ subfamm. \ n.} \end{array}$ 

Agnostidea see Stigmagnostus gen. n.

Agnostus exsulatus sp. n. p. 6 pl. i figs. 3-4 Cambrian (Middle) Argentina (Mendoza), Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Agnostus glabratus Angelin a synonym of Trinodus tardus [q.v.] p. 59, Kielan (1).

Agraulidae discussed p. 384, Diceratocephalinae referred p. 388, Kobayashi (4).

Agraulidae see Antagmopleura Lermontova [nom. nud.]

Agraulidae see Batenioides gen. n.

Agraulinae see Cyclolorenzella gen. n.

Agraulinae see Latilorenzella gen. n.

Agraulos divi Walcott type of Latilorenzella gen. n. to which is also referred A. regularis W. p. 390, Kobayashi (4).

Alacephalus gen. n. [family undetermined] for type A. contortus sp. n. p. 222 pl. xi figs. 11-13 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Alanisia migratoria n. sp. [nom. nud.] p. 82, Hupé (3).

Albertella pacifica Walcott referred to Chiawangella p. 354, Kobayashi (4).

Aldonaia Lermontova discussed, A. terea sp. n. with young stages p. 76 pl. v figs. 14-17, pl. vi figs. 1-10 text-figs. 25-26 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suverova Trud. paleont. Inst. S.S.R. 84 1960.

Aldonaidae discussed p. 75, Suvorova.

Aldoniidae see Planaspis gen. n.

Aldoniidae see Pumilina gen. n.

Alokistocarinae see Pedinocephalina.

Alsataspidae preferred to Selenecemidae and discussed p. 117, Whittard.

Americare gen. n. [Anomocaridae] for type Glyphaepis tetonensis Resser 1937 holotype figured p. 828 pl. c figs. 42-57 Cambrian (Middle) U.S.A. (Wyoming), Lochman & Hu J. Paleont. 34 5 1960.

Ameura go. p. 715 pl. xci fig. 3, Hoare & Burgess.

Amgaspidae fam. n. [Corynexochoidea] for Amgaspis Chernysheva 1956 p. 76, Chernysheva Principles of Palaeontology. Arthropods—Trilobitomorphs and Crustaceans 1960.

Amgaspis Chernysheva 1956 referred to Amgaspidae fam. n. p. 76, A. medius C. text-fig. 71, Chernysheva et al.

Amphilichas sp. p. 30, Bogdanov & Chugaeva.

Amphitryon radians (Barrande 1846) p. 65 pl. ii figs. 3-7, pl. iii fig. 11, A. sp. p. 67 pl. iii fig. 12, **Kielan** (1).

Ampyx depressus (Angelin) var. otradnica var. n. p. 32 [pl. i figs. 20-23] Ordovician (Middle), A. nordicus sp. n. p. 33 [pl. vi figs. 1-5], A. borealicus sp. n. p. 34 [pl. vi figs. 8-9] Ordovician (Upper) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Ampyx edgelli Reed type of Ampyxella gen. n. to which are possibly also referable A. latus McCoy, A. volborthi Schmidt and A. ? aculeatus Angelin p. 80, Dean (1),

Ampyx globifrons Olin a synonym of Raphiophorus gratus (Barrande) p. 167, A. tetragonus Angelin and A. t. var. gigas Linnarsson synonyms of Lonchodomas portlocki (B.) p. 171, Kielan (1).

Ampyx sp. p. 31, Bogdanov & Chugaeva.

Ampyx from Sahara p. 4406, Barlier et al.

Ampyxella gen. n. [Raphiophoridae] for type Ampyx edgelli Reed 1932 lectotype selected and figured p. 80 pl. xi figs. 1, 3-4, 6-7 Ordovician (Middle) Britain (Shropshire); A. latus McCoy, A. volborthi Schmidt, A. ? aculeatus Angelin possibly referable to Ampyzella p. 80, Dean Bull. Brit. Mus. (nat. Hist.) [Geol.] 4 4 1960.

Ampyzina taimyrica sp. n. p. 35 [pl. vi figs. 10-14] Ordovician (Middle) U.S.S.R. (Arctio), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Anchiops Delo referred to Monorakeidae (Carinopyginae) subfam. n. p. 8, Balashova (1).

Anechocephalus gen. n. [Ptychoparioidea (? Pterocephaliidae), may belong to Pterocephaliinae p. 84], A. trigranulatus sp. n. p. 92 pl. viii figs. 21-23 Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Anomocarella memor (Reed) p. 211 pl. xviii figs. 12-13, Repina.

Anomocaridae see Americare gen. n.

Anomocaridae see Kolbinella gen. n.

Anomocarioides—Coosia radugini Sivov referred p. 7, Ivshin.

Antagminae discussed p. 128, Suvorova.

Antagmopleura Lermontova [nom. nud.] [Agraulidae], A. sp. p. 202 pl. xvi figs. 13-14 Cambrian (Middle) U.S.S.R. (E. Sayan), Repina.

Antatlaeva n. gen. [nom. nud.] [Strenusevinse] p. 77, Hupé (3).

Aojia Resser & Endo 1937 a synonym of Lisania p. 368, Kobayashi (4).

Actiaspis gen. n. [Kainellinae] p. 246 for type A. oblonga sp. n. p. 247 pl. xiv figs. 7-13 text-fig. 6b, A. ovalis sp. n. p. 247 pl. xiv fig. 14 Ordovician (Lower) Korea (South), A. sp. indt. for Pseudo-kainella l b [8] sp. Kobayashi 1953 p. 248; A. lohanpoensis (Sheng) for Kainella l. S. p. 247 text-fig. 6e, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Apatokephalina Sivov discussed and tentatively referred to Richardsonellidae p. 242, Kobayashi (3).

Apatokephalina referred to Prospatokephalidae fam. n. p. 33, Rozova (2).

(Sib.

Apo
6d-f, 1

Apa kepha

fig. 14 Rozov

[196

Apa p. 237

Apa iam. n

Apo iv fig. Chern Apl for 2

Pap.

Api
n. p.
tumifi

N. ger Preho

Ar Ar syno

Sbor

A

A toke

p. 4 figs.

elucineli p. 2

vici of 195

257

30]

Coy, gelin

orus

and

type

and

(elbl

orthi able

nat.

-14]

lova

ino-

ero.

84], am-

ap.

figs.

red

auian

ae]

nia

6b,

do-

6c,

Apatokephaloides referred tentatively to Eurekiinae p. 237, Kobayashi (3).

Apatokephaloides referred to Proapatokephalidae fam. n. p. 33, Rozova (2).

Apatokephalus discussed and referred to Proapatokephalidae fam. n. p. 33, A. sibiricus sp. n. p. 35 pl. v fig. 14 Ordovician (Lower) U.S.S.R. (W. Siberia), Rozova Trud. Inst. geol. geophiz. AN S.S.S.R. (Sib. Otd.) 5 1960.

Apatokephalus hyotan Kobayashi p. 248 text-figs. 6d-f, Kobayashi (3).

Apatokephalus schoriensis Poletaeva [not seen] pl. iv fig. 22, A. subservatus P. [not seen] pl. iv fig. 23, Chernysheva et al.

Aphelaspidinae subfam. n. [Pterocephaliidae] p. 80 for Aphelaspis, Labiostria, Litocephalus, Taenora gen. n. and possibly Maladioidella, Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Aphelaspis referred to Aphelaspidinae subfam. n. p. 80, ventral cephalic sutures text-fig. 8E; "A." humifrons Resser may belong to Housiidae p. 74; N. gen. aff. A. of Palmer & Hazzard 1956 belongs to Prehousia gen. n. p. 78, Palmer (1).

Apianurus text-fig. 4, Whittington (2).

Aposolenopleura? brevifrons Rasetti see Canotaspis gen. n.

Arctinurus sp. pl. v fig. 21, Wilson & Majewske.

Arinichevia Polotaeva 1943 msc.—Olenina P. a synonym p. 26 n, Ivshin.

Aristoharpes taimyricus sp. n. p. 30 [pl. v figs. 20-21] Silurian (Lower) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Aristoharpes sp. p. 12 pl. i fig. 6 text-fig. 5d, Fillet (3).

Artokephalus Sivov & Egorova discussed p. 243, Kobayashi (3).

Artokephalus discussed and referred to Proapatokephalidae fam. n. p. 48, A. minimus Sivov with synonyms A. suburceus Egorova & A. squalaris S. p. 47 pl. v figs. 1-13, Rozova (2).

Asaphellus alimbeticus Balashova [nom. nud.] pl. vi figs. 3-5, Chernysheva et al.

Asaphellus tomkolensis Kobayashi tentatively excluding referred pygidium (1934 pl. 4 figs. 5-7) and including A. aff. gyracanthus and A. ? sp. of K. 1934 p. 259 pl. xiv figs. 15-24, Kobayashi (3),

Asaphid (indeterminate) p. 168, Dean (3).

Asaphidae Burmeister—revision of Middle Ordovician forms from Oslo with redeterminations of lists of Bragger 1884, 1887, Holtedahl 1909, Størmer 1953 p. 253, Henningsmoen.

Asaphidae—Upper Cambrian forms discussed p. 257, Kobayashi (3).

Asaphidae recorded p. 30, Bogdanov & Chugaeva.

Asaphidae gen. indét. p. 6, Pillet (3).

Asaphids (scraps of) p. 275, Stanton.

Asaphinae see Erdelia gen. n.

Asaphinae see Opsimasaphus gen. n.

Asaphinae see Protoptychopyge gen. n.

Asaphiscidae see Liopariinae subfam. n.

Asaphiscinae see Williamsina gen, n.

Asaphopsis nakamurai Kobayashi, A. cfr. n. p. 253, Kobayashi (3).

Asaphus elegantulus Gortani type of Dainellicauda gen. n. p. 253, Kobayashi (3).

Asaphus (Neoasaphus) heintzi sp. n. p. 231 pl. vii figs. 1-3, A. (N.) ludibundus Törnquist 1884 p. 233 pl. viii figs. 1-5, A. (N.) of. l. for A. n. sp. Holtedahl 1909 p. 234 pl. viii fig. 6, A. (N. l) sp. A p. 235 pl. xiv fig. 9, A. (N. l) sp. B p. 235 pl. xiv fig. 10 Ordovician (Middle) Norway (Oslo); A. debuchianus Brongniart 1817 a synonym of Ogygiocarie dilatata p. 217; Ogmasaphus may be a subgenus of A. p. 236, Henningsmoen Norsk geol. Tidsskr. 40 3-4 1960.

" Asaphus" luthardti table 1 p. 6, Volk.

Asaphus myops König referred to Dalmanites p. 280, Dean (4).

"Asaphus" nobilis Barrande restr. [lectotype chosen] discussed and generic attribution uncertain p. 76, A. n. B. 1852 partim (pl. 31 figs. 2-3, 6-8) and B. 1872 referred to Opsimasaphus jaanussoni sp. n. p. 79, "A" ingens B. discussed p. 77, A. i. of Olin 1906 referred to O. sp. p. 81, Kielan (1).

Asaphus priscus Lamansky type of Protoptychopyge gen. n. p. 125, Balashova (3).

Asioptychaspis Kobayashi discussed with list of referred spp. p. 400, Kobayashi (4).

Aspidagnostinae subfam. n. [Agnostidae] p. 60 for Aspidagnostus Whitehouse, Pokrovskaya Principles of Palaeontology. Arthropods—Trilobitomorphs and Crustaceans 1960.

Aspidagnostus Whitehouse 1936 referred to Aspidagnostinae subfam. n. p. 60, Chernysheva et al.

Astycoryphe gracilis (Barrande 1846) p. 211 pl. iii figs. 3-4, pl. iv fig. 3, Přibyl.

Astycoryphe ? junius (Billings 1869) p. 418 pl. liv figs. 11, 17-19, 22-23, 26-27, Whittington (1).

Atractopyge michelli (Reed) p. 126 pl. xvii figs. 12-16, Whittard.

Aulacopleura (A.) bohemica (Přibyl) for Cyphaspis sp. Agrell 1939 p. 105, Stubblefield.

Aulacopleura (Paraaulacopleura) halfari Strnad 1956 p. 343, Strnad (1). Aulacopleura (A.) konincki k. (Barrande) text-fig. 572, Müller.

Australosutura gen. n. [Brachymetopidae] p. 229 for type Cordania gardneri Mitchell 1922 lectotype selected and figured p. 230 pl. xxxix figs. 1-11, pl. xl figs. 1-20 text-fig. 2 (reconstr. showing crescentic rostral plate) Carboniferous Australia (N.S.W.) and Argentina (Chubut), Campbell & Goldring Palaeontology 3 2 1960.

Bailiella angusta sp. n. p. 373 pl. xix fig. 18 Cambrian (Middle) S. Korea, B. lantonoisi (Mansuy) with synonym B. ulrichi Resser & Endo immature forms described p. 374 pl. xxi figs. 1-10, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Baltagnostus hospitus sp. n. p. 7 pl. i figs. 5-6 [provisionally associated cephala may belong to Peronopsis ultima sp. n.], B. mendozensis sp. n. p. 8 pl. ifig. 7 Cambrian (Middle) Argentina (Mendoza), Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Baltagnostus wyomingensis sp. n. p. 822 pl. xeix figs. 1-4 Cambrian (Upper) U.S.A. (Wyoming), of. B. w. p. 822 pl. xeix fig. 32, Lochman & Hu J. Paleont. 34 5 1960.

Basidechenella see Dechenella.

Batenioides gen. n. [Agraulidae] p. 202 for type B. lermontovae sp. n. p. 203 pl. xv figs. 9-10 Cambrian (Lower) U.S.R. (E. Sayan), Repina Regional Strat. S.S.R. 4 1960.

Bathyurellus cf. tenuis Poulsen p. 451, Gobbett & Wilson.

Bathyurid[s] p. 451, Gobbett & Wilson.

Bathyurina sp. nov., f B. sp. nov. [both undescr.] p. 451, Gobbett & Wilson.

Bathyuriscellus Lermontova discussed and referred to Jakutinae p. 94, B. robustus L. lectotype selected p. 95 pl. viii figs. 3-9, pl. ix fig. 1, text-fig. 30, B. convexus sp. n. p. 98 pl. viii figs. 10-12 text-fig. 31, B. grandis sp. n. p. 100 pl. ix figs. 5-6 text-fig. 32, B. parous sp. n. p. 102 pl. ix fig. 2 text-fig. 33, B. viduus sp. n. p. 104 pl. ix figs. 3-4 text-fig. 34 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 34 1960.

Bathyuriscellus sp. p. 207 pl. viii fig. 6, Repina.

Bathyuriscopsis gen. n. [Jakutinae] p. 105 for type B. ultima sp. n. p. 106 pl. ix figs. 7-8 text-fig. 35 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Bergerolimbus n. gen. [nom. nud.] [Pseudoleninae] p. 77, B. elongatus, B. limbatissimus n. spp. [nom. nuda] p. 81, **Hupé** (3).

Bergeroniaspis divergens Lermontova p. 177 pl. v figs. 1-6, B. cf. d. pl. xii figs. 6-7; B. nitens and B. procera Suvorova spp. referred to Tungusella gen. n. p. 179, Repina.

Bergeroniaspis sp. pl. vi fig. 26, Suvorova.

Bergeroniellus spinosus Lerm. pl. xiv fig. 18 pars B, Suvorova.

Bergeronopsis n. gen. [nom. nud.] [Pseudoleninae] p. 77, B. zigzae n. sp. [nom. nud.] p. 81, Hupé (3),

Berkeia comes and B. nevadensis Resser 1942 synonyms of Kindbladia affinis p. 97, Palmer (1).

Bigotina Cobbold discussed with table of spp. p. 36, B. (Bigotina) C. diagnosis p. 37, B. (B.) angulata sp. n. p. 38 pl. ii figs. 20-21 text-fig. 10, B. (Bigotinella) subgen. n. p. 40 for type B. (Bigotinella) malykanica sp. n. p. 40 pl. ii figs. 22-25, pl. iii fig. 1, text-fig. 11 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Bigotina egregica sp. n. p. 217 pl. iii figs. 13-14, pl. vi fig. 10, B. comferica sp. n. p. 218 pl. vi figs. 11-13 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Bigotinella subgen, n. see Bigotina.

Bigotinops Hupé discussed p. 41, B. privus sp. n. p. 42 pl. iii fig. 2 text-fig. 12, B. patrius sp. n. p. 44 pl. iii figs. 3-4 text-fig. 13 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.R. 34 1960.

Bigotinops referred to Bigotinopsidae fam. n. p. 77, Hupé (3).

Bigotinopsidae fam. n. [no diagnosis] for Bigotinops, Pruvostina, Ouijjania, Tazemmourtia [nom. nud.] p. 77, Hupé Rep. XXI int. geol. Congr. 8 1960.

Binodaspis Lermontova discussed p. 128, B. prima L. 1951 [sp. n.—name provisionally suggested by L.] for B. sp. I L. p. 129 pl. xii figs. 7-8, 18, text-fig. 42, B. secunda sp. n. for B. sp. II L. p. 131 pl. xii figs. 9-12 text-fig. 43, B. paula sp. n. p. 133 pl. xii figs. 13-16 text-fig. 44 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Binodaspis spinosa Lermontova p. 201 pl. vi figs. 8-9, Repina.

Birmanitella subgen. n. see Birmanites.

Birmanites Sheng referred to Birmanitidae fam. n. p. 254, B. (Birmanitella) subgen. n. for type B. hupeiensis Yi p. 256 text-fig. 90 Ordovician (Upper) China, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Birmanitidae fam, n. [Dikelocephalidina] p. 254 for Birmanites and by implication Tropidopyge and Hagiorites, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Blackwelderia biloba Kobayashi referred to Damesops p. 353, B.? cornuta Endo to Diceratocephalus p. 391, Kobayashi (4).

Bojoscutellum Šnajdr 1958 discussed p. 98 (252), B. paliferum p. (Beyrich 1845) neotype chosen and figd. [excluding Barrande 1852 pl. 45 figs. 20-21 (to Breviscutellum inerme), pl. 45 fig. 4 and B. 1872, and with synonym Bronteus intermixtus Hawle & Corda lectotype chosen and figd.] p. 99 pl. x figs. 1-7, pl. xi figs. 1-4, pl. xii figs. 1, pl. xiii figs. 5-6, pl. xxxv fig. 14, text-figs. 6, 32-33, B. p. pseudopaliferum subsp. n. p. 103 (252) pl. xiv fig. 5, B. conspersum

(Pranti 1872 p. statum p. 105 35, ? I 5-6, pl Bohem

[1960

Bojo pl. liv

Bold Cambra assigna (pl. 5 Medd.

e figs.

Bol
[unde

Bol

Bol

Bon (Lowe 8.S.S

Bo Saya holot 49, \$

Br with Myst and Br

B

B for Bro xxx oho bohe ing

figs figd p. B. H. Ro

(lat

fro p. lec 5, η

(1). pp. ata la)

11

n),

na.

p.

p.

d

(Prantl 1949) holotype figd. including B. p. of B. 1872 p. 104 pl. xiii figs. 1-4 text-fig. 34, B. crassico-statum sp. n. including B. p. of B. 1852 pl. 45 fig. 4 p. 105 (253) pl. xiii fig. 7, pl. xiv figs. 1-4, text-fig. 35, f B. konéprusiense sp. n. p. 107 (253) pl. xi figs. 5-6, pl. xxxvi fig. 9, text-fig. 36 Devonian (Lower) Bohemia, šnajdr Rozpr. Úst. úst. geol. 26 1960.

Bojoscutellum ? pompilius (Billings 1869) p. 415 pl. liv figs. 13-16, 20-21, 24-25, Whittington (1).

Bolaspidella lucieae sp. n. p. 28 pl. ii figs. 16-17 Cambrian (Middle) Argentina (Mendoza); librigena assigned to Champlainia rectimargo Howell 1937 (pl. 5 fig. 34) referred to B. p. 29, Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Bolaspidella resseri (Miller) holotype figured pl. e figs. 36-38, Lochman & Hu.

Bolbocephalus of. seeleyi (Whitneld), B. sp. nov. [undescr.], ? B. p. 451, Gobbett & Wilson.

Bollandia soo Griffithides.

Bonnia veternosa sp. n. p. 199 pl. xi figs. 5-7, B. arguta sp. n. p. 200 pl. vii figs. 1-2 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.R. 4 1960.

Bonniella? hispanica Richter & Richter a synonym of Hicksia elvensis Delg. p. 82, Hupé (3).

Botomella Suvorova 1958 discussed with synonym Sayanella Pokrovskaya 1959 p. 149, B. ekaterinae S. holotype figured p. 150 pl. xiii figs. 9-12 text-fig. 49, Suvorova.

Brachymetopidae Prantl & Přibyl 1951 discussed with diagnosis and taken to include Cordania, Mystrocephala gen. n., Tschernyschewella, Piltonia and Brachymetopus p. 407, Whittington (1).

Brachymetopidae see Australosutura gen. n.

Brachymetopidae see Mystrocephala gen. n.

Brassicicephalus see Genus and species undet. 1.

Breviscutellum gen. n. [Scutelluidae] p. 172 (258) for type B. transversum (Hawle & Corda 1847) for Bronteus transversus H. & C. holotype figd, p. 173 pl. xxxii figs. 10-11, f 12, B. inerme (B. 1852) lectotype chosen and figd. [with synonyms Scutellum nováki and bohemicum Prantl & Přibyl holotype figd., and excluding records of i. by Kegel 1926 and Weber 1945 (latter to Decoroscutellum nov. sp.)] p. 174 pl. xxxiii figs. 1-8 text-fig. 54, B. ivanense (B. 1872) holotype figd. p. 177 pl. xxxiii figs. 5-6, B. obsoletum sp. n. p. 178 (259) pl. xxxiii figs. 10-11, pl. xxxiv fig. 1, B. angusticeps (B. 1846) with synonym B. sculptus H. & C. lectotypes of both chosen and figd. p. 178 pl. xxxiv figs. 2-7 Devonian (Lower) Bohemia, Snajdr Rozpr. Ost. úst. geol. 26 1960.

Broeggerolithus Lamont 1935 discussed with synonym Ulricholithus Bancroft p. 102, distinctions from Onnia p. 127, three groups of spp. recognised p. 103—(1) B. harnagensis (B.) for Salterolithus h. B. lectotype selected and figured p. 105 pl. xvi figs. 5, 9, 13, B. h. f. p. 106 pl. xvi figs. 1, B. cf. h. (B), for S. aff. h. B. 1949 p. 107 pl. xvi figs. 7, 11, B. f sp. p.

108 pl. xvi fig. 2; (2) B. ulrichi (B.) p. 109 pl. xvii figs. 10-11, B. broeggeri (B.) lectotype figured p. 111 pl. xvi figs. 3, 10, B. cf. b. (B.) p. 112 pl. xvi figs. 4, 8, 14, B. constrictus B. holotype figured p. 113 pl. xvi figs. 6, 12, B. soudleyensis (B.) including B. aff. s. Dean 1958 p. 115 pl. xvii figs. 1-9, 17, B. globiceps (B.) lectotype selected and figured p. 117 pl. xvii figs. 13, 15-16, B. sp. (? nov.) p. 119 pl. xvii fig. 12, B. simplex sp. n. [tentatively in group 2] p. 120 pl. xvii fig. 14; (3) B. longiceps (B.) lectotype selected and figured p. 121 pl. xviii figs. 1, 4-6, 10, B. transiens (B.) p. 123 pl. xviii figs. 2-3, 7-8, 11-14, B. cf. t. (B.) p. 126 pl. xviii fig. 9 Ordovician (Middle) Britain (Shropshire); Trinucleus discors Angelin may belong to B. p. 104, T. gibbifrons of Reed 1914 resembles B. g. p. 104, T. gibbifrons of Reed 1914 resembles B. sp. (? nov.) or B. simplex p. 121, Dean Bull. Brit. Mus. (nat. Hist.) [Geol.] 4 4 1960.

Broeggerolithus cf. broeggeri (Bancroft) p. 167, B. cf. globiceps (B.), B. longiceps (B.), B. cf. l. p. 168, Dean (3).

Brongniartella cf. parva Harper p. 167ff., B. bisulcata (M'Coy), B. ? sp. p. 168, Dean (3).

Brongniartella platynotus (Dalman 1828) with synonym Homalonotus inexpectatus Barrande 1872 p. 116 pl. xix figs. 1-3, Kielan (1).

Bronteus aff. costatus (Pusch), B. mirabilis N. Tchern, B. granulatus, B. sp. p. 121, Bel'skaya.

Bronteus hawlei Barrande (with synonym B. asperulus Hawle & Corda) type of Microscutellum gen. n. [q.v.] to which are also referred B. zippei and B. tenellus B. p. 92, B. pustulatus B. type of Metascutellum gen. n. [q.v.] p. 184, B. rhinoceros B. type of Cornuscutellum gen. n. p. 108, B. simulans R. type of Protoscutellum gen. n. [q.v.] p. 64, B. transversus H. & C. type of Breviscutellum gen. n. nermis and B. ivanensis B. p. 172, B. umbellifer Beyrich (with synonym B. sparsus H. & C.) type of Spiniscutellum gen. n. p. 111; B. aciculatus H. & C. nom. dub. p. 208, B. partschi B. referred to Kosovopeltis [q.v.] p. 70; B. carinatus and B. subtriangularis [q.v.] p. 70; B. carinatus and B. subtriangularis [q.v.] p. 70; B. carinatus and B. subtriangularis p. 100, B. berkeleyanus H. & C. of Platyscutellum formosum p. 119, B. sosia B. of P. spiniferum p. 129, B. gervilleicans B. of Poroscutellum infaustum p. 136, B. gigas H. & C. of Scabriscutellum caelebs p. 144, B. tardissimus B. of Scabriscutellum richteri p. 148, B. thysanopeltis B. of Thysanopeltis speciosa p. 157, B. scharyi B. of Kolhapeltis brevifrons p. 170, B. sculptus H. & C. of Paralejurus brongniarti p. 178, B. pulcher H. & C. of P. dormitzeri p. 196, Snajdr.

Bulaiaspis Lermontova discussed p. 184, B. prima L. p. 185 pl. xix figs. 1-2, B. taseevica sp. nov. [Repins 1956] holotype figured p. 186 pl. xix figs. 3-5, pl. ii figs. 7-12, B. peleduica sp. n. [nondeser. 1956] p. 188 pl. xix figs. 6-8, B. sajanica sp. n. p. 189 pl. iii figs. 1-4, B. timbata sp. n. p. 192 pl. iii figs. 5-8, B. mana sp. n. p. 193 pl. iii figs. 9-11, B. vologdini L. p. 184 pl. ii figs. 1-6, B. sp. pl. iii fig. 12, B. sp. pl. xix figs. 9-13 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.R. 4 1960.

Bulaiaspis Lermontova discussed with table of spp. p. 23, B. tolbatchanica sp. n. p. 24 pl. i figs. 5-9 text-figs. 3-4, B. modesta sp. n. with ontogenetic stages p. 27 pl. i figs. 10-18, pl. ii figs. 1-13, text-figs. 5-7, B. taseevica Repina 1956 p. 33 pl. ii figs. 14-19, text-figs. 8-9 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Bumastes [sic] barriensis (Murchison) p. 177, Squirrell & Tucker.

Bumastus sp. ? p. 270, Anderson.

Burnetia sp. cf. B. alta Resser p. 812 pl. xevi figs. 3-7, Lochman & Hu.

Bynumia see Genus and species undet. 3.

Bynumiella ? acuminata sp. n. p. 93 pl. x figs. 9-10 text-fig. 21 (reconstr.) Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Bynumina Resser 1942 discussed p. 93, B. globosa (Walcott) for Agraulos ? g. W. p. 94 pl. x fig. 8, Palmer (1).

Callavia choffati (Delg.)—C. lotzei Richter & Richter probably a synonym p. 82, Hupé (3).

Calliops cf. armatus Ulr. and Delo p. 30, Bogdanov & Chugaeva.

Calliops strasburgensis text-figs. 4c-d, Whittington (3).

Calliops taimyricus sp. n. p. 5 [pl. v figs. 9-12] Ordovioian (Upper) U.S.S.R. (Arctie), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Calvinella Walcott discussed with list of Asian spp. p. 405, "C." sp. indt. listed p. 336; C. sp. Kobayashi 1935 referred to Saukiella sp. p. 407, Kobayashi (4).

Calybole see Cyrtosymbole.

Calymenacea see Carmonidae fam. n.

Calymenacea see Synhomalonotidae fam. n.

Calymene (C.) beyeri R. & E. Richter, C. (C.) blumenbachi Brongn., C. sp. p. 350, Tomczykowa.

Calymene blumenbachii Brongniart, C. cf. beyeri (Richter), C. neointermedia (R. & E. R.), C. sp. p. 177, Squirrell & Tucker.

Calymene ex gr. blumenbachi Brongniart 1822 p. 70 pl. vii figs. 1-3, Maksimova.

Calymene collicephala Green text-figs. 512a-d, C. incerta Barrande text-fig. 589, Müller.

Calymene douvillei Mansuy type of Vietnamia gen. n. p. 43, Kobayashi (1).

Calymene hornýi n. sp. [nom. nud.] p. 114; C. tenera Barrande is apparently identical with Metacalymene baylei (B.), Vaněk.

Calymene (Colpocoryphe) inopinata, C. sp., C. pulchra table 1 p. 6, Volk.

Calymene nasuta Ulrich type of Spathacalymene gen. n. p. 892, Tillman.

Calymene neointermedia R. & E. Richter pp. 119, 123, Lawson.

Calymene sampelayoi sp. n. p. 136 pl. xiv figs. 1-2 Ordovician Spain (Ciudad Real), Linares Estud. geol. Inst. Mallada 9 17 1953.

Calymene sp. Whittard 1931 referred to Nescuretus grandior sp. n. p. 141, C. pompeckji Kobayashi referred to N. complanatus sp. n. and N. murchisoni (Salter) p. 150, Whittard.

"Calymene" sp. p. 115 pl. vii fig. 7, Kielan (1).

Calymenella Bergeron 1890—Echomalonotus Reed accepted provisionally as a synonym p. 162, Whittard.

Calymenella marocana n. sp., C. destombesi n. sp. [nom. nuda], C. cf. boisseli Berg., C. cf. bayani Trom. et Lebesc. p. 1538, **Destombes et al.** 

Calymenid indet. p. 167, Dean (3).

Calymenid pygidium p. 270, Anderson.

Calymenidae—Bohemian forms summarised p. 113, Vaněk.

Calymenidae—Endocrania referred p. 41, Synhomalonotidae separated off p. 42, Kobayashi (1).

Calymenidae—Pharostoma p. 133 and Neseuretus p. 141 referred, Whittard.

Calymenidae gen. u. sp. indet. table 1 p. 6, Volk,

Calymenidae see Mesoparia ord, n.

Calymenina referred to Ptychopariida p. 233, Kobayashi (3),

Calymeninae see Spathacalymene gen. n.

Camaraspis convexa (Whitfield) p. 813 pl. xovi figs. 8-17, Lochman & Hu.

Canotaspis gen. n. [Incertae sedis, compared with Aposolenopleura? brevifrons Rasetti] for type C. aliena sp. n. p. 29 pl. ii figs. 21-22 Cambrian (Middle) Argontina (Mendoza), Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Carbonocoryphe aff. bindemanni R. & E. Richter p. 271 pl. xii fig. 6, Prentice.

Carinopyge Maksimova 1955 discussed and referred to Carinopyginae subfam. n. p. 18, C. spinifera sp. n. p. 19 [pl. iv figs. 18 a-r] Ordovician (Middle) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Carinopyginae subfam. n. [Monorakeidae] p. 8 for Carinopyge and Anchiops, Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

p. 13

Ca.
n. p.

[19

Can

Car

for C

1960.

Ca

pp. 4

Ca sedis

Ce 14], Ordo [pl.

U.S. biosi

i. E

C

Kol

pl. pl. c. scu (Mi

pa

(3)

35 p. 30]

mene

pp.

geol.

retus

isoni

(1).

Reed

tard.

rom.

113,

etus

olk.

233,

kevi

with

dle)

hter

red

. n. 3.R.

rat.

for tat. Carmon Barrande 1872 referred to Carmonidae fam. n. p. 112, discussed and C. primus B. excluded pp. 44-47, C. mutilus (B. 1852) p. 112 pl. xvi fig. 4, pl. xvii figs. 1-6, text-figs. 5r, 6r, 30, 31, Kielan (1).

Carmonidae fam. n. [Calymenacea p. 47] p. 112 for Carmon Barrande, Kielan Paleont. polon. No. 11 1960.

Carnicia referred to Drevermanniinae subfam. n. p. 137, Chernysheva et al.

Caulaspis gen. n. p. 206 for type C. angustus sp. n. p. 207 pl. i figs. 7-8, C. latus sp. n. p. 209 pl. i figs. 5-6 Cambrian (Upper) U.S.S.R. (Salair), Rozova Trud. 1-nst. geol. geophix. AN SSSR (Sib. Otd.) 1 1960.

Caulaspis Rozova discussed and considered Incertae sedis p. 76, C. angustus R. holotype figd. p. 77 pl. iii fig. 16, C. latus R. holotype figd. p. 78 pl. iii fig. 12, Rozova (2).

Ceratevenkaspis parnaicus sp. n. p. 14 [pl. iv fig. 14], C. taimyricus sp. n. p. 15 [pl. iv figs. 9-11, 13] Ordovician (Upper), C. pentagonus sp. n. p. 14 [pl. iv figs. 7-86], C. borealicus [C. ? b. on pl. expl.] sp. n. p. 17 [pl. iv figs. 12 a-6] Ordovician (Middle) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Ceratocephala (C.) aff. vesiculosa (Beyr.) p. 350, Chlupáč & Vaněk.

Ceratocephala sp. table 1 p. 6, Volk.

Ceraurinella intermedia (Kielan 1955) for Ceraurus i. K. hypostome-described p. 127 pl. xxiii figs. 2-3 Kielan (1).

Ceraurinella cfr. mitis (Reed) p. 46 pl. v fig. 1, Kobayashi (1).

Ceraurinella text-figs. 1B, C, Whittington (2).

Ceraurinus nordicus sp. n. p. 22 [pl. iv fig. 19, pl. v fig. 2], C. borealicus sp. n. p. 23 [pl. iii fig. 31, pl. v fig. 22], C. taimyricus sp. n. p. 24 [pl. v figs. 3-4], C. confluens Barton p. 25 [pl. v fig. 1], C. of. tenusculptus Bradley p. 26 [pl. v fig. 6] Ordovician (Middle), C. kluevkaensis sp. n. p. 26 [pl. v fig. 7], C. icarus (Billings) p. 27 [pl. v figs. 23 a-6], C. marginatus Barton p. 28 [pl. v fig. 5] Ordovician (Upper) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Ceraurinus sp. p. 30f., Bogdanov & Chugaeva.

Ceraurus whittingtoni text-figs. la-h, Whittington (3).

Ceraurus sp. p. 30, Bogdanov & Chugaeva.

Ceraurus ? sp. p. 128 pl. xxi figs. 4-7 text-fig. 35; C. intermedius Kielan referred to Ceraurinella p. 127, Kielan (1).

Ceraurus text-fig. 1A, Whittington (2).

Cernuolimbus gen. n. [Pterocephaliinae] p. 84 for type C. orygmatos sp. n. p. 85 pl. viii figs. 1, 3, 5, 8, 11 text-fig. 18 (reconstr.), C. depressus sp. n. p. 85 pl. viii figs. 9-10, C. semigranulosus sp. n. p. 86 pl. viii figs. 2, 4, 6-7 Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Champlainia rectimargo Howell—assigned librigena (1937 pl. 5 fig. 34) referred to Bolaspidella p. 29, Poulsen (2),

Changia Sun discussed p. 401, Kobayashi (4).

Changshanocephalus Sun discussed and referred tentatively to Liopariinae subfam. n. p. 394, Kobayashi

Charchaqia discussed p. 257, Kobayashi (3).

Chariocephalus—assignment to Elviniidae uncertain p. 64, Palmer (1).

Chaunoproetus aff. carnicus (R. Richter) p. 178 pl. xxix figs. 8-9, Selwood.

Cheilocephalus sp. undet. p. 815 pl. xevii figs. 11-13, Lochman & Hu.

Cheilocephalus spp. p. 94 pl. x figs. 13-14, Palmer

Cheiropyge Diener excluded from Brachymetopidae p. 408, Whittington (1),

Cheiruridae see Hammatocneminae subfam. n.

Cheirurinae see Emsurella gen. n.

Cheirurinae see Eocheirurus gen. n.

Cheirurus cf. C. insignis Beyrich p. 268, Churkin & Langenheim.

Cheirurus insocialis of Barrande 1872 [non 1852] referred to Actinopeltis barrandei sp. n. p. 130, C. neglectus of B. 1872 partim [pl. 3 fig. 18] probably belongs to A. insocialis p.140, Kielan (1).

Cheirurus sp. pl. v fig. 20, Wilson & Majewske.

Chiawangella Chu 1959 [not seen] discussed p. 353, C. pustuloca C. text-fig. 2j, Albertella pacifica Walcott referred to C. p. 354, Kobayashi (4).

Chondragraulos minussensis Lermontova p. 201 pl. xvi figs. 4-5, C. ujaricus sp. n. p. 201 pl. xiv figs. 10-12, C. sp. p. 202 pl. viii fig. 9 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Chondranomocare irbinica sp. n. p. 209 pl. xvi figs. 8-9, pl. xviii figs. 1-3 Cambrian (Middle) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Chuangia Walcott 1911 discussion of referred spp. 361, C. taihakuensis Kobayashi listed p. 333, C. sp. p. 334; C. nais var. persicum King type of Iranochuangia gen. n. p. 363, C. n. of K. referred to I. narghunensis sp. n. p. 363, Kobayashi (4).

Chuangioides Chu 1959 [not seen] discussed p. 360, C. punctatus C. text-fig. 3 i, Kobayashi (4).

Chuangiopsis Sivov discussed and referred to Pagodiinae p. 370, Kobayashi (4).

Clavagnostinae subfam. n. [Agnostidae] for Clavagnostus Howell p. 59, Pokrovskaya Principles of Palaeontology. Arthropods—Trilobitomorphs and Crustaceans 1960.

Clavagnostus Howell 1937 referred to Clavagnostinae subfam. n. p. 59, C. repandus (Westergård) pl. i figs. 12-13, Chernysheva et al.

Clavagnostus chipiqueneis (Rusconi 1952) for Culipagnostus c. R. [reinterpreted as a pygidium] p. 9 pl. i fig. 14; C. calensis R. 1950 excluded from C. p. 10, Poulsen (2).

Cliffia lataegenae (Wilson) p. 814 pl. xev figs.1-7,

Cliffia latagenae see Genus and species undetermined 3.

Coenaspis see Pedinocephalina gen. n.

Coignouina see Otarion.

Colpocoryphe 800 Calymene.

Colpocoryphe cf. aragoi Rouault p. 1537 ff., Destombes et al.

Colpocoryphe bohemica n. sp. [nom. nud.] p. 113, Vaněk.

Colpocoryphe grandis see Paracolpocoryphe.

Comluella linnarssoni (Kiaer) from Morocco p. 77, Hupé (3).

Conokephaluella n. gen. [nom. nud.] [Ellipsostre-nuellinae] p. 77, Hupé (3).

Coosella see Genus et sp. ind. (pygidium).

Coosia see Genus et sp. ind. (librigena).

Coosia radugini Sivov referred to Anomocarioides p. 7, Ivshin.

Coosina ariston (Walcott) p. 816 pl. xevii figs. 15-20, cf. C. sp. undet. p. 818 pl. xevii figs. 29-31, Lochman & Hu.

Cordania Clarke 1892 discussed with diagnosis (including presence of facial sutures) p. 408, C. cyclurus (Hall & Clarke 1888) lectotype selected and figured p. 409 pl. li figs. 1-7, 9-10, 12-13, 19, C. falcata sp. n. Devonian (Lower) U.S.A. (Oklahoma) p. 411 pl. li figs. 8, 11, 14-18, pl. lii figs. 3, 5, C. macrobius (Billings 1869) p. 412 pl. liii figs. 1-14, 17-18, 20-21, text-fig. 1b, C. becraftensis C. 1900 lectotype selected and figured with synonym C. hudsonica C. (holotype figured) p. 412 pl. lii figs. 6-9, 11, 13-14, 16-19, C. gasepiou C. 1908 holotype figured p. 413 pl. lii figs. 1-2, C. sp. p. 413 pl. lii figs. 4; C. gardneri Mitchell 1922 excluded [by R. Goldring] from C. p. 407, C. pulchra Cooper & Cloud 1938 type of Mystrocephala gen. n. p. 413 to which is also referred C. rara Stumm, Whittington J. Paleont. 34 3 1960.

Cordania gardneri Mitchell type of Australosutura gen, n, p. 229, Campbell & Goldbing in Amos et al.

Cornovica gen. n. [Encrinuridae—forming a group with Dindymene and Plasiaspie] p. 122 for type C. didymograpti sp. n. for Dindymene sp. Whittard 1953 p. 123 pl. xvii figs. 1-11 Ordovician (Middle) Britain (Shropshire), Whittard Palaeontogr. Soc. [Monogr.] 113 (for 1959) 1960.

Cornuproetus see Proetus.

Cornuproctus (C.) aff. buchi Hawle & Corda p. 160, Cavet.

Cornuproetus cf. holzapfeli (Novák 1890) p. 49 pl. iii figs. 1, 5, pl. v fig. 12, C. altaicus sp. n. p. 50 pl. ii figs. 1-4, pl. v fig. 11 Devonian (Middle) U.S.S.R. (Altai), Maksimova Devonian and Carboniferous trilobites of the Rudny Altai 1960.

Cornuscutellum gen. n. [Scutelluidae] p. 108 (253) for type Bronteus rhinoceros Barrande 1872 lectotype chosen and figd. p. 109 pl. xviii figs. 2-7, pl. xix figs. 1-2, text-figs. 37-38 Devonian (Lower) Bohemia, Snajdr Rozpr. Úst. úst. geol. 26 1960.

Coronura ? spec. p. 129 pl. i fig. 3, Bürgl.

Corynexochoidea see Amgaspidae fam. n.

Corynexochoidea—Triniidae referred p. 78, Chernysheva et al.

Costonia Whittard 1956 discussed p. 86, C. ultima (Bancroft) cast of holotype figured p. 87 pl. xii figs. 1-12, C. elegans sp. n. for C. sp. nov. Dean 1958 p. 89 pl. xiii figs. 1-6, 8-9 Ordovician (Middle) Britain (Shropshire), Dean Bull. Brit. Mus. (nat. Hist.) [Geol.] 4 4 1960.

Crepicephalus granulosus Hall & Whitfield type of Elburgia gen. n. p. 68, Palmer (1).

Cromus 800 Encrinurus.

Crotalocephalus myops F. Roemer 1850 p. 71 pl. vii figs. 4-9, C. aff. uratjubensis Weber 1932 p. 74 pl. vii fig. 11, C. sp. (hypostome) p. 75 pl. vii fig. 10, Maksimova.

Cryphops see Phacops.

Cryptolithus grenieri Berg. p. 1538, Destombes et al.

Cryptolithus cf. portlocki of Størmer 1930 probably referable to Onnia [q.v.] p. 138, C. cf. gibbifrons of Whittington 1938 may be Broeggerolithus sp. (? nov.) or B. simplex sp. n. p. 121, Dean (1).

Cryptolithus text-figs. 3D-F, Whittington (2).

Culipagnostus chipiquensis Rusconi reinterpreted as a pygidium and referred to Clavagnostus [q.v.] p.9, Poulsen (2).

Cummingella sp. aff. balladoolensis Reed 1942 for Griffithides cf. minor of Frech 1906 and aff. m. of Rakusz 1932 p. 38 pl. i fig. 1-2, Bouček & Přibyl.

C<sub>3</sub>
125,
C<sub>3</sub>
Lore

Boge

[1

Cybei Whi

Bpp. tonk
Ross
yent
Chir
Uni

of I (pyg p. 8 p. 8 (Up polo

C

also

to 0

(A.

C

exciand A.:

(

I. 1 Phi 271

Ric (Go Th of. nep pl. C.

prop. E. R.

al.

Ĉ

lle)

oe.

60,

R.

pe

17-

58 sin st.)

of

10.

al.

v.)

.9.

Cybele Lovén 1845—type taken as C. bellatula and Cybelina Reed thus an objective synonym p. 125, Whittard.

Cybele aff. planifrons Web., C. sp. indet. p. 30, Bogdanov & Chugaeva.

Cybelina Reed an objective synonym of Cybele p. 125, Whittard.

Cyclolorenzella gen. n. [Agraulinae] p. 389 for type Lorenzella quadrata Kobayashi text-fig. 9 j, following spp. also referred acalle and armata Walcott, tonkinensis Mansuy, rotundata, conveza, ogurai Resser & Endo, parabola Lu, pustulosa, subcylindrica, yentaiensis, kushanensis Chu Cambrian (Middle) China and Korea, Kobayashi J. Fao. Sci. Tokyo Univ. [2] 12 2 1960.

Cyclopyge gallica Tjernvik p. 217, Tjernvik.

Cyclopyge prisca (Barrande) eye text-fig. 505a [see also Pricyclopyge], Müller.

Oyclopyge quadrangularis sp. n. p. 83 including C. rediviva of Nicholson & Etheridge 1878-80 and of Reed 1904 p. 83 pl. ix figs. 1-9 text-figs. 22, 23 (pygidial ontogeny), C. gigantea (Barrande 1872) p. 86 pl. x fig. 2, C. sp. a p. 87 pl. x fig. 7, C. sp. b p. 87 pl. x fig. 3, C. sp. c p. 88 pl. x fig. 1 Ordovician (Upper) Poland (Lysa Gora) etc., Kielan Palaeont. polon. No. 11 1960.

Cyphaspides sp. p. 350, Chlupáč & Vaněk.

Cyphaspides sp. Strnad 1956 [recte 1957] referred to Otarion (O.) couvinianum sp. n. p. 340, Strnad (1).

Cyphaspis novella Barrande referred to Harpidella p. 199 (219), Přibyl.

Cyphaspis sp. Agrell 1939 referred to Aulacopleura (A.) bohemica (Přibyl) p. 105, Stubblefield.

Cyphoproetus rugosus see Prionopeltis sp.

Cyrtometopus? decacanthus Angelin restr. [cephalon excluded] lectotype selected [thorax and pygidium] and referred to Diacanthaspis p. 103, C. octacanthus A. restr. [non fig. 6] lectotype selected and referred to Pseudosphaerexochus p. 134, Kielan (1).

Oyrtoproetus Reed 1943 discussed with diagnosis p. 19, C. dobsinensis (Illés 1902) for Griffithides d. I. holotype figured p. 21 pl. i figs. 3-4, Bouček & Přibyl.

Cyrtosymbole (Waribole) cf. aequalis (Meyer) p. 271 pl. xii fig. 5, Prentice.

Cyrtosymbole (Waribole) aff. conifera R. & E. Richter p. 179 pl. xxix fig. 7, C. (W.) aff. italica (Gortani) p. 179 pl. xxix fig. 6, C. (? W.) dunhevedensis Thomas nom. correct. type not found p. 180, C. (W.) of. warsteinensis R. & E. R. p. 156, C. (? Calybole) cf. nepic R. & E. R. p. 180, C. (? Macrobole) aff. [no aff. on pl. expl.] bergica R. Richter p. 180 pl. xxix fig. 5, C. (M.) drewerensis R. & E. R. sens. nov. with probable synonym C. (M.) d. longieuta R. & E. R. p. 180 pl. xxix figs. 14-16, C. (M.) duodecimae R. & E. R. p. 181 pl. xxix figs. 17, C. (M.) aff. blax R. & E. R. p. 182 pl. xxix figs. 12-13, C. (M.) sp. p. 183 pl. xxix figs. 10-11, Selwood.

Cyrtosymbole l sp. p. 54 pl. ix figs. 5, 12 pars, Maksimova.

Cyrtosymbolidae see Drevermanniinae subfam. n.

Custispina see Phillibole.

Daguinaspis ambroggii Hupé & Abadie meraspis showing undivided palpebral lobes text-fig. 2, Hupé (2).

Dainellicauda gen. n. [Dikelokephalininae] p. 253 for type Asaphus elegantulus Gortani text-fig. 7c Ordovician (Middle) Himalayas, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Dalmanites longicaudatus orientalis Reed 1906 p. 9 pl. i figs. 4-5, 10 pars a [sic], text-fig. 5a, Pillet (3).

Dalmanites myops (König) for Asaphus m. K. 1825 with synonym D. vulgaris (Salter) of which selected lectotype is holotype of D. m. p. 280, Dean (4).

Dalmanites oklahomae Richardson referred to Neoprobolium and age considered Devonian (Lower) p. 53, Frederickson.

Dalmanites vulgaris (Salter) p. 117, Lawson.

Dalmanites vulgaris (Salter), D. v. var. bituberculata Reed p. 177, Squirrell & Tucker.

Dalmanites sp. Tschernyschew 1893 referred to Proetus sp. p. 47, Maksimova.

Dalmanitina mucronata (Brongniart 1822) with synonyms D. kiaeri Troedsson and D. proeva grandis (Barrande) p. 118 pl. xix fig. 4, pl. xx figs. 5-11, D. olini Temple 1952 p. 119 pl. xix figs. 5-6, pl. xx figs. 12-13, D. o. or D. m. [young stages] p. 120 pl. xix figs. 7-9, pl. xx figs. 1-4, D. m. and D. o. migration p. 39 ff., Kielan (1).

Dalmanitina proaeva of Bordet et al. 1959 referred to Pterygometopus (P.) dagon p. 5n, Pillet (3).

Dalmanitina sp. text-fig. 514 pars, D. (D.) socialis (Barrande) text-fig. 596, Müller.

Dalmanitina sp. p. 1538, Destombes et al.

? Dalmanitina sp. p. 334, Cave.

Damesella paronai Airaghi listed p. 334; D. nitida Resser & Endo referred to Paramenocephalites gen. n. p. 381, Kobayashi (4).

Damesellidae discussed p. 350 [see also Paramenomoniinae subfam. n.], Kobayashi (4).

Damesops Chu 1959 [not seen] discussed p. 352, D. convexus C. text-fig. 2e, D. bilobus (Kobayashi) for Blackwelderia b. K. 1942 text-fig. 2f, Kobayashi (4).

Daryagnostus [sic] incertus (Brögger) p. 10, Noakes et al.

Dawsonia sibirica Lazarenko mse. [nom. nud.] pl. ii figs. 24-25, Chernysheva et al.

Dechenella (Basidechenella) dombrowiensis Guerich 1896 p. 339 pl. i figs. 1-5, pl. iv fig. 18, Strnad (1).

Dechenella minima sp. n. p. 136 pl. i figs. 6a-c Devonian (Middle) Japan (Iwate), Okubo Earth Science [Chikyū Kagaku] No. 4 1951.

Deckera sp. cf. D. completa Wilson p. 812 pl. xovi fig. 34, Lochman & Hu.

Decoroscutellum Šnajdr 1958 discussed p. 79 (249), D. (Decoroscutellum) discussed p. 80 (249), D. (D.) haidingeri (Barrande 1846) lectotype chosen and figd. p. 80 pl. v figs. 10-15, pl. vi figs. 1-7, pl. xxxvi fig. 11, text-fig. 26, D. (D.) binotatum (B. 1872) holotype figd. p. 84 pl. vi figs. 8-11, pl. vii fig. 1, text-fig. 27, D. (D.) lepidum l. (Bouček 1933) holotype figd. p. 85 pl. vii figs. 2-6 text-fig. 28, D. (D.) texoptatum subsp. n. p. 87 (250) pl. vii fig. 9 text-fig. 29, D. (D.) ascriptium sp. n. p. 88 (250) pl. ix fig. 5, D. (D.) uralicum (Weber 1945) lectotype chosen and figd. p. 89 (250) pl. xxxvi figs. 5-6, D. (Fleziscutellum) subgen. n. for type D. (F.) hanuši sp. n. including Scutellum umbelliferum of Prantl & Přibyl 1947 partim (pl. 6 fig. 2) p. 90 (251) pl. vi fig. 12, pl. vii figs. 7-8, text-fig. 30 Silurian (Upper) Bohemia, D. nov. sp. for Bronteus angusticeps inermis of Weber 1945 (ast. god. 26 1960.

Delgadella caudata (Delgado) from Morocco p. 77,

Dellea suada (Walcott) p. 813 pl. xcvi figs. 22-27, Lochman & Hu.

Denckmannites see Phacopidella.

Denemarkia frontalis (Hawle & Corda 1847) lectotype selected and immature forms illustrated p. 188 (212) pl. i figs. 1-10, Přibyl.

Diacanthaspis decacantha (Angelin 1854) restr. for Cyrtometopus ? d. A. lectotype [thorax and pygidium] selected and figured p. 103 pl. xv figs. 1-3, pl. xvi figs. 2a-b, 3, pl. xvii figs. 7-8 text-fig. 27, Kielan (1).

Dianops see Phacops.

Diceratocephalinae referred to Agraulidae p. 391, Kobayashi (4).

Diceratocephalus Lu discussed and Blackwelderia? cornuta Endo referred p. 391, Kobayashi (4).

Dicranopeltoides cf. D. decipiens (Weller) p. 268, Churkin & Langenheim.

Dictyites dictys (Walcott) p. 334, D. depressa Kobayashi p. 336, Kobayashi (4).

Dikellocephalus multicinctus Hall & Whitfield type of Hederacauda gen. n. p. 250, Kobayashi (3).

Dikelocephalidae see Girvanopyge gen. n.

Dikelocephalidina see Birmanitidae fam. n.

Dikelocephalinae discussed p. 249, Kobayashi (3).

Dikelocephalinae see Hederacauda gen. n.

Dikelocephalites flabelliformis Sun may belong to Pterocephaliidae p. 84, Palmer (1).

Dikelocephalopsis Poletaeva discussed and referred to Dikelokephalininae p. 252, D. amsassensis P. text-figs. 8 a-b, Kobayashi (3).

Dikelokephalina parva sp. n. p. 252 pl. xiii figs. 27-30, D. conica sp. n. p. 253 pl. xiii fig. 31 Ordovician (Lower) Korea (South), Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Dikelokephalininae discussed and taken to include Dikelokephalina, Asaphopsis, Hungioides, Dactylocephalus, Asaphopsoides, Dainellicauda gen. n. and probably Leimitzia, Dikelocephalopsis and Temnurá p. 251, Kobayashi (3).

Dikelokephalininae see Dainellicauda gen. n.

Dindymene Hawle & Corda 1847 discussed with list of spp. p. 146, D. fridericiaugusti H. & C. p. 148 pl. xxviii figs. 1-2, text-fig. 9b, D. ornata Linnarsson 1869 [may be a synonym of D. heidingeri Barrande p. 147] lectotype chosen and refigured with synonym D. spinulosa Olin 1906 refigured p. 148 pl. xxvi fig. 6, l. xxvii figs. 3-5, pl. xxii fig. 5, pl. xxvii figs. 3-5, pl. xxii fig. 5, pl. xxvii fig. 4, pl. xxx figs. 1-3, text-fig. 43 Ordovician (Upper) Poland and Sweden, D. plasi sp. n. p. 151 pl. xxix figs. 1-3 text-fig. 42 Ordovician (Middle) Bohemia; D. pulchra Olin type of Eodingmen gen. n. p. 144, D. cornuta O. a nom. dub. D. venusta O. a synonym of Raphiophorus gratus (Barrande), D. minuta Reed 1931 a nom. dub. and may be part of a Trinodus p. 147, Kielan Palaeont. polon. No. 11 1960.

Dindymene Hawle & Corda 1847 diagnosis p. 121, considered to form with Plasiaspis and Cornovica gen. n. a compact group of Encrinuridae, D. sp. Whittard 1953 referred to C. didymograpti sp. n. p. 123, Whittard.

Dindymeninae subfam. nov. [recte Přibyl 1953] for Dindymene and Eodindymene gen. n. pp. 52, 143, Plasiaspis excluded p. 144, Kielan (1).

Dindymeninae see Eodindymene gen. n.

Dinesidae Lermontova discussed with synonym Proerbidae Hupé p. 139, Suvorova.

Dinesidae see Tumulina gen. n.

Dionide subrotundata sp. n. p. 161 pl. xxxi fig. 4, pl. xxxii figs. 1-3, pl. xxxiii fig. 7, pl. xxxv fig. 5, text-fig. 46, D. decorata sp. n. p. 163 pl. xxxi figs. 1-3 text-fig. 47 Ordovician (Upper) Scandinavia and Poland, Kielan Palaeont. polon. No. 11 1960.

Diplagnostus jarillensis Rusconi 1952 / referred cephalon p. 10 pl. i fig. 9, Poulsen (2).

Diplapatokephalus referred to Proapatokephalidae fam. n. p. 33, Rozova (2).

Dokimocephalidae—Acrocephalitinae referred p. 383, Kobayashi (4),

grego

[19

D

D

Resenyment with and E. a. E. p. 3

D

Kob

refer

fam D p. 1 Pal

Cru

D D D D D

to E
D.
orpl
figs.
[excepted]

syn p. p. p. D. 28-(Ne

fig. berg (H. K. spec hall

(H. Pa)

Kie

Ki

to

an

ci.

de

nd

ra

th

48

on

m

6,

t.

D.

m

us

d t.

ca

m

d

Dokimocephalus Walcott discussed, D. pernasutus (W.) holotype figured with possible synonym D. gregori W. p. 95 pl. xi figs. 18, 21 [sic], Palmer (1).

Dokimocephalus sp. p. 813 pl. xevi fig. 37, Lochman & Hu.

Dolerolenidae see Sajanaspis gen. n.

Doryagnostus see Daryagnostus.

Dorypyge richthofeni Dames discussed with several Resser & Endo 1937 and Resser 1942 spp. as synonyms p. 347, D. r. laevis Walcott holotype figured with synonyms p. 348 pl. xix figs. 19-25, laiwuensis and manchuriensis Kobayashi and perconveza R. & E. also subspp. of r. p. 347, D. ? grandispinosa R. & E. p. 349, D. taihakuensis and D. ofr. t. K. listed p. 334, Kobayashi (4).

Dorypygidae discussed p. 343, Pseudokoldinioidia referred p. 349, Kobayashi (4).

Drepanura premesnili Bergeron listed p. 333, Kobayashi (4).

Drevermannia referred to Drevermanniinae subfam. n. p. 137, Chernysheva et al.

Drevermanniinae subfam. n. [Cyrtosymbolidae] p. 137 for Drevermannia, Eodrevermannia, Carnicia, Palpebralia, Formonia, Maksimova Principles of Palaeontology. Arthropods—Trilobitomorphs and Crustaceans 1960.

Dreyfussina see Phacopidina.

Drumaspis—assignment to Elviniidae uncertain p. 64, Palmer (1).

Dulanaspis Chugaeva excluded from Scutelluidae p. 53, Šnajdr.

Dunderbergia Walcott 1924 discussed and referred to Elviniidae p. 65, ventral cephalic sutures text-fig. 8 c, D. bigranulosa sp. n. with possible librigenal dimorphism and a pathological pygidium p. 66 pl. v figs. 10-13, 15-23, D. nitida (Hall & Whitfield) [excluding pygidium of Walcott 1925 pl. 16 figs. 6-7 to Housia] lectotype selected and figured with synonym D. simulator (H. & W.) holotype figured p. 67 pl. iv figs. 14-21, 23-24, D. polybothra sp. n. 67 pl. v figs. 14-21, 23-24, D. polybothra sp. n. D. variagranula Palmer p. 68 pl. iv figs. 22, 25, 28-29, D. v. ? pl. iv fig. 26 Cambrian (Upper) U.S.A. (Nevada), D. quadrata Kobayashi [of which D. v. may be a synonym] holotype figured p. 68 pl. iv fig. 27; D. maculosa (H. & W.) [type of Megadunder-bergia] holotype figured pl. v fig. 5, D. pustulosa (H. & W.) holotype figured pl. v fig. 8, D. canadensis K. and D. convexa K. should all be restricted to type specimens and considered indeterminate p. 66, D. halli Resser referred to Housia p. 75, D. granulosa (H. & W.) type of Elburgia gen. n. p. 68, Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Dysplanus see Illaenus.

Ectillaenus cf. leptopleura (Holm) listed p. 30, Kielan (1).

Ectillaenus sarkaensis (Novak) text-fig. 560B, Müller.

Edelsteinaspidae discussed p. 123, Suvorova.

Edelsteinaspidae—Laticephalus Pokrovskaya referred p. 209, Repina.

Edelsteinaspis gracilis Lerm. MS pl. xiv fig. 18 pars 6, Suvorova.

Edelsteinaspis ornatus Lermontova p. 208 pl. viii figs. 10-11, pl. xiv figs. 7-9, Repina.

Eilura Resser & Endo discussed and referred to Menocephalitinae subfam. n. p. 382; E. quadrata E. excluded, Kobayashi (4).

Elasmaspis gen. n. [Monorakeinae] p. 1030 for type E. speciosa sp. n. p. 1031 pl. figs. 1 (I-V) Ordovician (Upper) U.S.S.R. (Siberia), Kramarenko Doklady Akad. Nauk SSSR (ns) 109 5 1956.

Elasmaspis Kramarenko referred to Monorakeidae (Elasmaspisinae) subfam. n. p. 7, Balashova (1).

Elasmaspisinae subtam. n. [Monorakeidae] p. 7 for Elasmaspis Kramarenko, Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Elatelimbus gen. n. [Acrocephalitidae] for type E. gammatus sp. n. p. 213 pl. xviii figs. 9-11 Cambrian (Middle) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.R. 4 1960.

Elburgia gen. n. [Elviniidae] p. 68 for type Crepicephalus granulosus Hall & Whitfield 1877 holotype refigured p. 69 pl. vi figs. 16-17, 19, E. quinnensis (Resser) for Taenicephalus q. R. holotype refigured p. 69 pl. vi figs. 11-13, 15, text-fig. 10 (reconstr.) Cambrian (Upper) U.S.A. (Nevada); Elvinia sp. of Palmer & Hazzard 1956 belongs to Elburgia, Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1980.

Elganellus Suvorova discussed p. 46, E. probus S. p. 47 pl. iii figs. 5-7 text-fig. 14, E. pensus S. p. 48 pl. iii figs. 8-16 text-figs. 15-16, E. acceptus S. p. 52 pl. iii figs. 17-18, pl. iv figs. 1-5, text-figs. 17-18, e. elegans S. with young stages p. 55 pl. iv figs. 6-19, pl. v figs. 1-13, text-figs. 19-20, Suvorova.

Ellipsocephalidae discussed p. 61, Suvorova.

Ellipsocephalinae discussed p. 63, Suvorova.

Ellipsocephalus hoffi (v. Schlotheim) text-fig. 531, Müller.

Ellipsocephalus uniformis n. sp. [nom. nud.], E. sp. p. 81, **Hupé** (3).

Ellipsuella n. gen. [nom. nud.] [Ellipsostrenuellinae] p. 77, E. tangarfae, E. kautzkii n. spp. [nom. nuda] p. 81, Hupé (3).

Elvinia Walcott 1924 discussed, E. roemeri (Shumard) with all Reser's 1937, 1938, 1942 spp. of E. [except 2 following] as synonyms p. 70 pl. vi fig. 7 text-fig. 11 (reconstr.), E. granulata R. holotype figured with synonym E. ruedemanni R. p. 71 pl. vi fig. 4, E. hamburgensis (R.) for Parairvingella h. R. (holotype figured) may be a synonym of E. g. p. 71 pl. vi fig. 5, E. ? sp. p. 71 pl. vi fig. 6; E. sp. of Palmer & Hazzard 1956 referred to Elburgia gen. n. p. 69, Palmer (1).

Elvinia roemeri (Shumard) with immature stages p. 814 pl. xevi figs. 38-47, Lochman & Hu.

Elviniella gen. n. [Elviniidae] p. 71 for type E. laevis sp. n. p. 72 pl. vi figs. 9-10, E. sp. [may belong to E. l.] pl. vi figs. 8, 14 Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Elviniidae Kobayashi discussed and taken to include Elvinia, Elviniella and Elburgia genn. n., Dunderbergia, Irvingella, relations discussed p. 64, Palmer (1).

Elviniidae see Elburgia gen. n.

Elviniidae see Elviniella gen. n.

Elviniidae see Hypostomes A-E.

Emsurella gen. n. [Cheirurinae] for type E. humila [humilla on pl. expl.] sp. n. p. 28 pl. iii figs. 7-9, E. ? laevigata sp. n. p. 30 pl. iii figs. 6, 10-11 Cambrian (Upper) U.S.S.R. (Salair), Rozova Trud. Inst. geol. geophiz. AN SSSR (Sib. Otd.) 5 1960.

Emsurina Sivov discussed p. 24, E. minuta sp. n. p. 25 pl. iii figs. 1-2, E. fulita sp. n. p. 26 pl. iii figs. 3-5 Cambrian (Upper) U.S.S.R. (Salair), Rozova Trud. Inst. geol. geophiz. AN SSSR (Sib. Otd.) 5

Encrinuridae discussed p. 120; Dindymene, Plasiaspis and Cornovica gen. n. form a compact group p. 121, Whittard.

Encrinuridae see Cornovica gen. n.

Encrinurus (Cromus) beaumonti (Barrande) p. 268, Churkin & Langenheim.

Encrinurus punctatus Emmrich text-fig. 587, Müller.

Encrinurus punctatus (Wahlenburg), E. cf. variolaris Brongniart, E. sp. p. 177, Squirrell & Tucker.

Encrinurus sp. p. 38, McKerrow & Campbell.

Encrinurus sp. nov. [undeser.] p. 119, Lawson.

Endocrania Kobayashi referred to Calymenidae and compared with Ptychometopus p. 41, Kobayashi (1).

Endymionia kazachstanica Balashova [not seen] pl. iv fig. 6, Chernysheva et al.

Entomostracites granulatus Wahlenberg 1818 partim (pl. 2 fig. 4\*) refigured and referred to Leonaspis olini [q.v.] p. 101 [see also Tretaspis granulata], Kielan (1).

Eobronteus Reed 1928 discussed p. 53 (245), E. reedi Sinclair type of Protobronteus gen. n. p. 55, Snajdr.

Eccheirurus gen. n. [Cheirurinae] p. 16 for type E. salairicus sp. n. p. 18 pl. i figs. 14-16, 18-19, E. subtilis sp. n. p. 20 pl. i fig. 17, pl. ii figs. 1-3, E. probus sp. n. p. 22 pl. ii figs. 4-9 Cambrian (Upper) U.S.S.R. (Salair), Rozova Trud. Inst. geol. geophiz. AN SSSR (Sib. Otd.) 5 1960.

Eochuangiinae discussed p. 359, Kobayashi (4).

Eccryphops see Phacops.

Eodindymene gen. n. [Dindymeninae] p. 144 for type Dindymene pulchra Olin 1906 lectotype selected and refigured including D. sp. of O. partin (pl. 1 fig. 26) refigured p. 144 pl. xxvii figs. 1-2, pl. xxviii figs. 3-4, pl. xxix fig. 6, text-figs. 9a, 40 Ordovician (Upper) Sweden and Poland, Kielan Palaeont. polon. No. 11 1960.

Eodiscid f [called Serrodiscus f by C. J. Stubblefield p. 105] p. 103, Bassett in Bassett & Walton.

Eodrevermannia referred to Drevermanniinae subfam. n. p. 137, Chernysheva et al.

Eohomalonotid ? pp. 141, 168, Cavet.

Echomalonotus Reed 1918 accepted provisionally as a synonym of Calymenella p. 162, Whittard.

Eorobergia sp. p. 30, Bogdanov & Chugaeva.

Eosaukia Lu discussed p. 404, Kobayashi (4).

Erbia Lermontova discussed p. 141, E. granulosa g. L. lectotype selected p. 143 pl. xiii figs. 1-3 texf-fig. 46, E. g. astricts subsp. n. p. 145 pl. xiii figs. 4-8 text-fig. 47 Cambrian (Middle) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Erbia granulosa Lermontova p. 204 pl. xv figs. 7-8, E. sibirica (Schmidt) p. 204 pl. viii figs. 12-13, pl. ix fig. 7, pl. xv fig. 4, pl. xvi figs. 6-7, pl. xvii figs. 12-13, Repina.

Erdelia gen. n. [Asaphinae] (nom. n. for Maja Lomovitskaya 1955) for type M. subsulcata L. 1955 p. 145, E. venusta (L.) pl. iii fig. 6 Tremadoc U.S.S.R. (W. Siberia), Lomovitskaya Principles of Palaeontology. Arthropods—Trilobitomorphs and Crustaceans 1960.

Eremiproetus see Proetus.

Eremiproetus cf. eremita (Barrande 1852) for Proetus sp. Tschernyschew 1893 refigured p. 53 pl. i fig. 8, Maksimova.

Euloma kelleri Balashova [nom. nud.] pl. x fig. E. kazachstanica B. [nom. nud.] pl. x fig. 5, Chernysheva et al.

Euloma new species [like E. geinitzi] p. 43, Lochman-Balk.

Eurekiinae Hupé may belong to Monkaspidae, Apatokephaloides tentatively included p. 237, Kobayashi (3).

Eurudagnostus referred to Geragnostidae p. 234, Kobayashi (3).

Evenkaspis norilskiensis sp. n. p. 13 [pl. iv figs. 15-17] Ordovician (Middle) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Eye (Kobe E. con (Midd Univ.

[19

Fle planis Fle declin

declin Vanël Fle

Fo gen.

n. p.

Ga
pygic
Poul

Ge

Ge Cam Toky

Ge Hom is tr

Ge

Gam Sei.

G

(1). G pele Ball

type 7e C J. F

1933 [om

o. I

gen

4).

for

ed

viii

an

on.

ble

b-

lly

1).

Ú.

an

R.

8,

3,

R.

t-

4,

Eymekops Resser & Endo discussed, E. expansus (Kobayashi) for Lioparia e. K. p. 395 pl. xix fig. 17, E. carinata sp. n. p. 396 pl. xxi fig. 11 [sio] Cambrian (Middle) S. Kores, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Flexicalymene cambrensis (Salter) p. 157, F. planimarginata (Reed) p. 158, Whittard.

Flexicalymene incerta (Barr. 1846) pl. x and F. declinata (Hawle & Corda 1847) so referred p. 113, Vaněk.

Flexicalymene sp. p. 334, Cave.

Flexiscutellum subgen. n. see Decoroscutellum.

Forbesia brevifrons Angelin referred to Ogmocnemis gen. n. p. 69, Kielan (1).

Formonia referred to Drevermanniinae subfam. n. p. 137, Chernysheva et al.

Gallagnostus solitariensis Rusconi reinterpreted as a pygidium and referred to Phoidagnostus [q.v.] p. 14, Poulsen (2).

Gentilaspis n. gen. [nom. nud.] [Olenellid] p. 76, Hupé (3).

Gentilaspis tuberosissima Hupé [nom. nud.] text-fig. 4B, Hupé (2).

Geragnostella cambria sp. n. p. 342 pl. xix fig. 8 Cambrian (Upper) S. Korea, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Geragnostidae discussed p. 233, Rudagnostus and Eurudagnostus referred p. 234, Kobayashi (3).

Geragnostus considered generically separable from Homagnostus to which G. tumidosus (Hall & Whitfield) is transferred p. 62, Palmer (1).

Geragnostus obsoletus sp. n. p. 342 pl. xxi figs. 16-17 Cambrian (Upper) Korea (South), Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Geragnostus wimani Tjernvik p. 217, Tjernvik.

Geragnostus sp. p. 58 pl. i fig. 4 text-fig. 12, Kielan (1),

Geragnostus (Micragnostus) [sp.] straddling Trempeleau/Tremadoc junction in Montana p. 45, Lochman-Balk.

Girvanopyge gen. n. [? Dikelocephalidae] p. 254 for type Lichapyge ? problematica Reed 1906 text-fig. 7e Ordovician (Upper) Britain (Scotland), Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Glaphyraspis Resser 1937—Raaschella Loohman 1938 a synonym p. 279, RASETTI in Harrington et al. [omitted Z.R. 96 11].

Glaphyraspis occidentalis Lochman for Raaschella o. L. p. 815 pl. xovii figs. 1-8, Lochman & Hu.

Chyphaspis tetonensis Resser type of Americane gen. n. p. 828, Lochman & Hu.

Glyptagnostus reticulatus r. (Angelin) and G. r. angelini (Resser) stratigraphic positions in U. S. A., G. n. sp. [undescr.] p. 1941, Palmer (2).

Goycoia Rusconi 1950 discussed and type designated G. tellecheai R. 1950 p. 30 pl. ii fig. 24, G. brevicauda sp. n. p. 31 pl. ii fig. 23, G. pecoralis sp. n. p. 32 pl. ii fig. 25 Cambrian (Middle) Argentina (Mendoza), Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Grandioculus sp. p. 334, Kobayashi (4).

Granularia Poletayeva diagnosis, G. protōlenorum Lermontova 1951 neotype selected p. 162 pl. xiv figs. 17-18, pl. xv, pl. xvi, text-fig. 54, G. bidjensis P. [nom. nud.] p. 167, Suvorova.

Granularia obrutchevi Poletaeva p. 209 pl. xiv figs. 1-6, Repina.

Granulariidae discussed p. 162, Suvorova.

Griffithides Portlock 1843 discussed p. 30, G. (Bollandia) Reed 1943 so referred and discussed p. 30, G. (B.) rozlozeniki Rakusz 1932 lectotype selected and figured p. 31 pl. vi figs. 1-3; G. dobsinensis Illés referred to Cyrtoproctus p. 21, G. cf. and aff. minor see Cummingella, Bouček & Přibyl.

Griffithides verrucosus Gemmellaro referred to Neoproetus p. 4, Ruggieri.

Grönwallia may belong to Pterocephaliidae p. 80, Palmer (1).

Grossia subgen. n. see Leonaspis.

Hagiorites Kobayashi discussed with implied reference to Birmanitidae fam. n. p. 256, Kobayashi (3).

Hammatoeneminae subfam. n. [Cheiruridae] for Hammatoenemis gen. n. p. 141, discussed p. 48, Kielan Palaeont. polon. No. 11 1960.

Hammatocnemis gen. n. [Cheiruridae (Hammatocneminae)] p. 141 (affinities discussed p. 48) for type H. tetrasulcatus sp. n. p. 141 pl. xxv fig. 3, pl. xxvi figs. 2-4, pl. xxvii figs. 6-8, text-figs. 38, 39 (ontogeny of cranidium) Ordovician (Upper) Poland (Lysa Gora), Kielan Palaeont. polon. No. 11 1960.

Haniwa quadrata Kobayashi p. 335, Kobayashi

Harpes (H.) radians r. Reinh. Richter 1863 p. 110, H. (H.) r. subsp. indet. pl. xii fig. 9, H. (H.) r. asimus Schwarzbach 1950 p. 110, Alberti.

Harpes reticulatus Hawle & Corda 1847 lectotype chosen p. 24 pl. vi figs. 5a-6, H. r. var. altaica Weber holotype figured p. 26 pl. i figs. 1a-6, Maksimova.

Harpes reticulatus Hawle & Corda p. 350, Chlupáč & Vaněk.

Harpes ruderalis Hawle & Corda text-fig. 574, H. ungula Sternberg text-fig. 575, Müller.

Harpes sp. p. 31, Bogdanov & Chugaeva.

Harpes sp. p. 350, Tomezykowa.

[1

I

subf

3-4 Reg

I

I

Kin

Kol

refe

Kol

K.

R. ]

figu

I

Chu

xiv

Am

Hu

Mo

ma figs

figs

Jal

Ch

inc

poi Su

Ja

fig J.

Tr

p. 3

Harpes sp. indet. p. 335, Strnad (1).

Harpidella McCoy 1849 nov. emend. discussed p. 218, H. novella (Barrande 1852) for Cyphaspis n. B. p. 219 pl. iii figs. 5-6, H. megalops (McCoy) with possible synonym H. elegantula (Lovén), H. christyi (Hall), H. girardeauensis (Shumard) all so referred p. 219, Přibyl.

Hawleia see Placoparia.

Hebediscus lermontovae sp. n. p. 174 pl. vii figs. 4-6 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Hederacauda gen. n. [Dikelocephalinae] for type Dikellocephalus multicinctus Hall & Whitfield p. 250 text-figs. 8 c-d Cambrian (Upper) U.S.A., Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Hemigyraspis desiderata, H. sp. table 1 p. 6, Volk.

Hibbertia sanctacrucensis sp. n. p. 157 pl. xxxiv figs. 4-6, pl. xxxv fig. 8, text-fig. 44 Ordovician (Upper) Poland (Lysa Gora), H. sp. p. 159 pl. xxxii fig. 7, Kielan Palaeont. polon. No. 11 1960.

Hicksia elvensis Delg.—Bonniella ? hispanica Richter & Richter belongs p. 82, Hupé (3).

Hintzeia glabella sp. n. p. 262 pl. xiii fig. 32 Ordovician (Lower) Korea (South), Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Holcacephalus tenerus (Walcott) with immature stages including protaspides p. 823 pl. xeviii figs. 1-52, Lochman & Hu.

Holmiinae see Sinskia gen. n.

Homagnostus Howell discussed and considered generically separable from Agnostus and Geragnostus p. 62, H. tumidosus (Hall & Whitfield) transferred from G. p. 63 pl. iv figs. 1-2, H. obesus (Belt) p. 63 pl. iv figs. 7-9; H. acutus Kobayashi referred with query to Pseudagnostus p. 62, Palmer (1).

Homalonotidae see Mesoparia ord. n.

Homalonotus barratti Woodward may be a Phacopid p. 104, Stubblefield.

Homalonotus inexpectatus Barrande a synonym of Brongniartella platynotus (Dalman) p. 118, Kielan (1).

Homalonotus knightii König p. 120, Lawson.

Homalonotus knightii König p. 177, Squirrell & Tucker.

Homalonotus sp. p. 350, Tomczykowa.

Housia Walcott discussed p. 74, incomplete separation of last thoracic segment from pygidium described, ventral cephalic sutures text-fig. 8 n, H. halli (Resser) for Dunderbergia h. R. holotype figured p. 75 pl. vii fig. 8, H. ovata sp. n, with transitory pygidia p. 75 pl. vii figs. 1-7, 9 text-fig. 12 (reconstr.) Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960

Housia sp. p. 814 pl. xevi figs. 1-2, Lochman &

Housiidae discussed and taken to include Housia, Prehousia and Parahousia genn. n., and possibly Morosa gen. n., "Aphelaspis" tumifrons Resser, Acrocephalaspis and Ullaspis p. 74, Palmer (1).

Housiidae see Morosa gen. n.

Housiidae see Parahousia and Prehousia genn. n.

Howelluella n. gen. [nom. nud.] [Strenuellinae] p. 77, **Hupé** (3).

Hsiaoshia Endo discussed p. 408, Kobayashi (4).

Hukasawaia Kobayashi discussed p. 244, H. cylindrica K. text-fig. 6 g; Portentosus is probably allied to H. p. 243, Kobayashi (3).

Hungioides graphicus, ? H. table 1 p. 6, Volk.

Hystricurinae discussed p. 382, Kobayashi (4).

Hystricurus megalops Kobayashi 1934 p. 235 pl. xiii fig. 20, H. ofr. m. p. 236 pl. xiii fig. 21, H. calvus sp. n. p. 236 pl. xiv figs. 5-6 Ordovician (Lower) Korea (South), Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Hystricurus wilsoni sp. n. for H. aff. flectimembrus of Hallam 1958 p. 457 pl. xv figs. 1-14 text-fig. 6 (reconstr.), Ordovician (Lower) Spitsbergen, Gobbett Geol. Mag. 97 6 1960.

Hystricurus sp. p. 451, Gobbett & Wilson.

Iddingsia Walcott discussed with relations to Plataspella analysed p. 95, I. robusta (W.) holotype figured with synonym I. nevadensis Resser holotype figured p. 96 pl. xi figs. 13-16, I. similis (W.), I. utahensis R., I. missouriensis R. with synonym I. crassimarginata R. diagnoses; I. s. of Bell et al. 1952 partim (pl. 31 fig. 2) referred to Plataspella p. 96, I. ? quinnensis R. referred to Sigmocheilus gen. n. p. 90, Palmer (1).

Illaenide indét. p. 1537, Destombes et al.

Illaenopsis ? acuticaudata Hicks referred to Seleneceme p. 118, Whittard.

"Illaenus" angelini Holm 1882 pl. xiv fig. 6, "I." cf. a. H. p. 95 pl. xiv figs. 7-9, "I." sp. p. 95 pl. xiv fig. 5, Kielan (1).

Illaenus glaberrimus (Hawle & Corda 1847) considered nom. dub. p. 209, **Šnajdr.** 

Illaenus lacertus with muscle scars text-figs. 4g, h, Whittington (3).

Illaenus cf. latiaxiatus Raym., I. aff. marginalis R., I. sp. p. 30, Bogdanov & Chugaeva.

Illaenus loretzi, I. sp. table 1 p. 6, Volk.

Illaenus (Dysplanus) namshimensis Reed 1915 p. 11 pl. i figs. 7, 10 pars b [sic], text-figs. 2, 5c, Pillet (3).

Illaenus sp. p. 1539, Destombes et al.

0]

1 &

ibly

).

. n.

ase

(4).

H.

k.

pl.

vus ver)

niv.

g. 6 bett

to

ype

I.

ym

al.

ella

to

95

on-

h,

ilis

3).

Illaenus sp. p. 177, Squirrell & Tucker.

Inouyella discussed and referred to Liopariinae subfam. n. p. 391, Kobayashi (4).

Inouyina subquadratica sp. n. p. 215 pl. iv figs. 3-4 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Inouyops Resser 1942 a synonym of Paragraulos p. 385, Kobayashi (4).

Iranochuangia gen. n. [Leiostegiinae] p. 363 for type I. persicum for Chuangia nais var. persicum King 1937 text-fig. 3k, I. narghunensis sp. n. for C. n. of K. 1930 p. 363 Cambrian (Upper) Iran, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Irvingella Ulrich & Resser 1924 discussed and referred to Elviniidae p. 72, I. (Parairvingella) Kobayashia recognisable taxon, I. (P.) angustilimbatus K. holotype figured with synonym P. intermedia R. p. 73 pl. vi figs. 2-3, I. (P.) eurekeneis R. holotype figured p. 74 pl. vi fig. 1, Palmer (1).

Isotelus gigas text-fig. 4b, Whittington (3).

Isotelus ef. maximus Locke p. 30, Bogdanov & Chugaeva.

Isotelus cf. platyrhachis Steinhardt 1874 p. 251 pl. xiv figs. 5-7, Henningsmoen.

Isotelus sp. large articulated specimen p. 18 fig. 1, Amsden & Ham.

Issafenia n. gen. [nom. nud.] [Olenellid] p. 76, Hupé (3).

Ithyektyphus Shaw 1956 discussed and relations to Modocia analysed p. 829, I. tetonensis (Miller) type material discussed and holotype figured p. 830 pl. c figs. 15-27, I. nevadensis (Palmer) for M. n. P. p. 830, I. tipperaryensis (M.) holotype figured pl. c figs. 1-3, Lochman & Hu.

Iwayaspis nom. nud. p. 257, Kobayashi (3).

Jakutidae Suvorova diagnosis and division into Jakutinae and Malykaninae subfam. n. p. 84, Suvorova.

Jakutidae fam. nov. [recte Suvorova 1959] p. 72, Chernysheva et al.

Jakutidae see Manaspis gen. n.

Jakutinae subfam. nov. [Suvorova] taken to include Jakutus, Bathyuriscellus, Judaiella, Parapoliella, Lenaspis and Bathyuriscopsis gen. n. p. 86, Suvorova.

Jakutus Lermontova discussed and referred to Jakutinae, J. mirus sp. n. p. 86 pl. viii figs. 1-2 textfig. 27 Cambrian (Lower) U.S.S.R. (Siberian Platform), J. fuzovi Galimova [not seen] listed p. 86, Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Josina gen. n. [Menomoniidae] p. 830 for type J. festiva sp. n. p. 831 pl. c figs. 28-35 Cambrian (Middle) U.S.A. (Wyoming), Lochman & Hu J. Paleont. 34 5 1960.

Judaiella Lermontova referred to Jakutinae p. 86, Suvorova.

Judaiella inepta sp. n. p. 206 pl. i figs. 1-3 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Kabutocrania gen. n. [Tingocephalinae] p. 358 for type K. fossula sp. n. p. 359 pl. xx fig. 19 text-fig. 2i Cambrian (Upper) S. Korea, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Kainella euryrachis Kobayashi p. 245 "text-fig. 6h; K. lohanpoensis Sheng referred to Aotiaspis gen. n. p. 247, Kobayashi (3).

Kainellina Poletayeva 1945 [not seen] [Remopleurididae] diagnosis p. 116, K. kousnezkiensis P. pl. iv figs. 27-28, Chernysheva et al.

Kainellinae see Aotiaspis gen. n.

Kameschkowiella [Kameschkoviella on pl. and pl. expl.] gen. n. [Protolenidae] for type K. zhuruvlevae sp. n. p. 176 pl. xi figs. 8-10 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.R. 4 1960.

Kaolishania Sun 1924 list of spp., K. granulosa Kobayashi 1933 p. 355 pl. xx figs. 8-12, pl. xxi fig. 14, K. ? latiura sp. n. for aff. K. ? orientalis of K. 1935 p. 356 pl. xxi fig. 19 Cambrian (Upper) S. Kores, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

? Kaolishaniella transita Sun p. 356 pl. xx figs. 13-16, Kobayashi (4).

Kaolishaniidae discussed and classified into Kaolishaniinae and Tingocephalinae p. 354, Kobayashi (4).

Kendallina eryon (Hall) p. 811 pl. xev figs. 24-30, Lochman & Hu.

Kettneraspis see Leonaspis.

Kindbladia Frederickson 1948 discussed, K. affinis (Walcott) holotype figured with synonyms Berkeia nevadensis (holotype figured) and B. comes Resser p. 97 pl. xi figs. 17, 19-20, Palmer (1).

Kingstonella subgen. n. see Kingstonia.

Kingstonia Walcott discussed p. 66 and classified into K. (Kingstonia) and K. (Kingstonella) [subgen. n.] for type K. walcotti Resser p. 70; K. gibbosa Lermontova, K. globosa, K. paichinensiformis, K. salairica Sivov spp. referred to Parakoldinia gen. n. p. 52, Rozova Trud. Inst. geol. geophiz. AN S.S.S.R. (Sib. Otd.) 5 1960.

Kingstonia see Genus and species undet. 3.

Kingstoninae discussed p. 61 ff., Rozova (2).

Kjerulfia adaioides, K. serksi n. spp. [nom. nuda] p. 81, **Hupé** (3).

Kokuria discussed and referred to Liopariinae subfam. n. p. 391, Kobayashi (4).

Kolbinella gen. n. [Anomocaridae] for type K. prima sp. n. p. 210 pl. iv figs. 1-2 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1980

Koldiniella Sivov discussed and may belong to Tsinaniidae p. 397, Kobayashi (4).

Koldinioidea [sic] sp. p. 335, Kobayashi (4).

Kolihapeltis Prantl & Přibyl 1947 discussed p. 163 (257), K. parabolina (Barrande 1882) lectotype chosen and figd. with synonym Scutellum (K.) lanceolatum P. & P. holotype figd. p. 165 pl. xxxi figs. 6-11 text-figs. 51, 52, K. perlonga (B. 1872) holotype figd. p. 167 pl. xxxi figs. 5, K. linguada (Novák 1884) holotype figd. p. 168 pl. xxxi figs. 12-13, K. chlupáći sp. n. for S. (K.) l. of P. & P. 1947 figd. Devonian Czechoslovakia (Bohemia) p. 169 (258) pl. xxxi fig. 14, K. brevifrons (B. 1852) with synonym Bronteus scharyi B. lectotypes of both chosen and figd. p. 170 pl. xxxii figs. 1-4 text-fig. 53, K. sp. p. 172 pl. xxxi fig. 15, Snajdr Rozpr. Ust. úst. geol. 26 1960.

Komaspidae discussed and Irvingella removed p. 73, Palmer (1).

Komaspidae see Terechtaspis gen. n.

Kootenia abacanica Poletaeva p. 199 pl. x figs. 10-11, K. sp. p. 199 pl. xv figs. 1-3, Repina.

Kooteniella slatkowskii (Schmidt) p. 198 pl. ix figs. 4-6, pl. xv figs. 11-12, pl. xvi figs. 1-3, K. sp. p. 199 pl. viii figs. 14-15, K. cranidium pl. ix fig. 1 pars, Repins.

Koptura ? sp. p. 212 pl. xviii fig. 8, Repina.

Koraipsis spinus Kobayashi p. 262 pl. xiii figs. 16a-b text-fig. 11, Protopliomerops punctatus K. may belong to K. p. 262, Kobayashi (3).

Kormagnostus Resser 1938 discussed, K. ? propinquus sp. n. p. 11 pl. i fig. 10 Cambrian (Middle) Argentina (Mendoza), Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Kormagnostus simplex Resser p. 822 pl. xcix figs. 5-31, K. sp. undet. p. 799, Lochman & Hu.

Kosovopeltis Šnajdr 1958 discussed p. 65 (247), K. svobodai Š. [holotype figured] with immature stages p. 66 pl. ii figs. 3-6, pl. iii figs. 1-16, 18-26, pl. iv figs. 1-2, text-figs. 21-22, K. partschi p. Barrande 1846) restr. lectotype chosen and figured with synonyms Bronteus subtriangularis and B. carinatus Hawle & Corda [both lectotypes chosen and figd.] & including B. simulans of B. 1852 partim (pl. 48 figs. 31-32) p. 70 pl. iv figs. 3-12 text-fig. 23, K. partschi inopinata subsp. n. including B. p. of B. 1852 (pl. 46 fig. 31) p. 73 (247) pl. iv figs. 13-16, K. parvispina sp. n. p. 73 (248) pl. iv figs. 13-14 text-fig. 24, K. hornji sp. n. including B. p. of B. 1852 (pl. 46 figs. 23-24, 125) p. 75 (248) pl. v figs. 2-7 text-fig. 25, K. nuntia (B. 1852) holotype figd. p. 76 pl. v fig. 8 Silurian (Upper) Bohemia, K. tchernychevae sp. n. for B. of. p. of Weber 1945 figd. p. 78 (249) pl. xxxvi fig. 7 Silurian U.S.S.R. (Urals), Snajdr Rozpr. Ust. ust. gool. 26 1960.

Krolina gen. n. [Protolenidae] p. 175 for type K. pressulata 59. n. p. 176 pl. i figs. 4-6 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Kushanopyge serrata Chu 1959 discussed and may be referable to Monkaspis p. 269 text-fig. 12c, Kobayashi (3).

Kyphocephalus sp. p. 813 pl. xevi figs. 35-36, Lochman & Hu.

Labiostria Palmer 1954 restricted to type and referred to Aphelaspidinae subtam. n. p. 80; L. platifrons P. tentatively referred to Taenora gen. n. p. 84, L. sigmoidalis P. removed from L. and related to Sigmocheilus gen. n. p. 80, Pterocephalina texana Resser 1942 probably referable to L. p. 90, Palmer (1).

Laticephalus Pokrovskaya referred to Edelsteinaspidae, L. sp. p. 209 pl. vii fig. 10, Repina.

Latilorenzella gen. n. [Agraulinae] p. 390 for type Agraulos divi Walcott text-fig. 9g, also including A. regularis W. Cambrian (Middle) China, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Lecanopygidae see Minicephalus gen. n.

Leimitzia Sdzuy discussed and referred to Dikelokephalininae p. 251, Kobayashi (3).

Leiostegijnae discussed p. 359, Kobayashi (4).

Leiostegiinae see Iranochuangia gen. n.

Lenaspis Suvorova discussed and referred to Jakutinae p. 88, L. opipara S. holotype figured p. 89 pl. xi figs. 20-22 text-fig. 28, L. limata S. holotype figured p. 91 pl. xii fig. 1 text-fig. 29, Superveys.

Leonaspis (Grossia) subgen. n. p. 115 for type Acidaspis myops Reinh. Richter 1863 p. 116 pl. xii figs. 7-8 Devonian (Middle) Germany (Thüringia), Alberti Mitt. geol. Staatsinst. Hamb. 29 1960.

Leonaspis coronata (Salter), L. sp. p. 177, Squirrell & Tucker.

Leonaspis (Acanthalomina) minuta (Barrande) p. 269, Churkin & Langenheim.

Leonaspis (Kettneraspis) moravica (Strnad 1956) p. 344 pl. ii figs. 6-9, Strnad (1).

Leonaspis olini Troedsson 1918 refigured including Entomostracites granulatus Wahlenberg 1818 partim (pl. 2 fig. 4\*) refigured and with synonym Acanthaloma mirka Marek 1952 p. 98 pl. vii figs. 5-6, pl. xiv figs. 1-4, pl. xv fig. 4, pl. xviii fig. 6 text-fig. 25, L. centrina (Dalman 1828) lectotype selected and figured p. 101 pl. xv fig. 5 text-fig. 26, Kielan (1).

Leonaspis sp. p. 116, Lawson.

Lermontoviella Pokrovskaya referred to Ptychopariidae, L. porrecta 5p. n. p. 201 pl. xiii figs. 9-10 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.R. 4 1960.

Lie proba b, Ko Lie referr

[196

Koba Lic gen.

p. p

Lie Ordo

Lie plour Li

type Ogyg pl. x and r. of fig. Gore

No.

Li
Kobi
p. 39
sp. 1
Koro
p. 39
1960

Li Yok Inon ? O J. H

Aoj spp. 191 Chi: 2 19

> Suv Suv Ito (He

1

pl. p. (U) 960]

r type mbrian

Strat

d may

. 12e,

35-36,

e and 30; L.

en. n.

texana

almer

einas-

type

ayashi

ikelo-

d to

gured

ta S.

. 29,

type

6 pl.

ngia),

irrell

e) p.

1956)

ding artim

loma

. xiv

and (1).

reho-

9-10 pins

1).

Liaoningaspis taitzehoensis Chu 1959 discussed and probably referable to Monkaspis p. 268 text-figs. 12a-b, Kobayashi (3).

Liaotungia Reeser & Endo 1937 discussed and referred to Menocephalitinae subfam. n. p. 381, Kobayashi (4).

Lichapyge? problematica Reed type of Girvanopyge gen. n. p. 254, Kobayashi (3).

Lichas cf. anglicus Beyer, L. hiroutus Fletcher, L. sp. p. 177, Squirrell & Tucker.

Lichas araneus Lindström 1885—request to ICZN for ruling whether L. aranea Holzapfel 1895 is a homonym p. 233, Tripp.

Lichengia Kobayashi discussed and referred to Ordosiinae p. 372, Kobayashi (4).

Lichida taken to include Lichina and Odonto-pleurina p. 233, Kobayashi (3).

Liobole see Phillibole.

Liocnemis gen. n. [Pterygometopinae] p. 120 for type Phacops recursus Linnarsson 1869 with synonym Ogygia ? apiculata L. 1866 p. 121 pl. ix figs. 11-12, pl. xxii figs. 8-11, pl. xxii figs. 1-2, text-fig. 32 Sweden and Poland, L. concinnus sp. n. for Pterygometopus r. of Kielan 1956 partim (pl. 2 figs. 2-3) p. 123 pl. ix fig. 10, pl. xxii figs. 3-7, text-fig. 33 Poland (Lysa Gora) Ordovician (Upper), Kielan Palaeont. polon. No. 11 1960.

Lioparia Lorenz discussed with synonym Yokusenia Kobayashi and referred to Liopariinae subfam. n. p. 393, L. latelimbata (Dames) discussed, L. conicula sp. n. p. 394 pl. xxi fig. 15 Cambrian (Upper) S. Korea; L. ezpansus Kobayashi referred to Eymekops p. 395, Kobayashi J. Fao. Sci. Tokyo Univ. [2] 12 2 1960.

Liopariinae subfam, n. [replacement name for Yokuseninae] [Asaphisoidae] p. 391 for Lioparia, Inouyella, Kokuria, ? Mapania, ? Changshanocephalus, ? Orloviella spp. [q.v.], ? Meticaspina, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Lisania Walcott 1911 discussed with synonym Aojia Resser & Endo, discussion and list of referred spp. p. 368, L. convergens sp. n. for L. alala of W. 1913 partim (pl. 15 fig. 19a) p. 369 Cambrian (Middle) China, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 1960.

Litaspis gen. n. [Solenopleurellinae] for type L. pudens sp. n. p. 137 pl. xii figs. 17, 19, text-fig. 45 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.R. 84 1960.

Litocephalus Resser 1937 discussed and referred to Aphelaspidinae subfam. n. p. 81, L. bilobatus (Hall & Whitfield) p. 82 pl. vii figs. 24-27 text-fig. 16 (reconstr.), L. granulomarginatus sp. n. p. 82 pl. viii figs. 14, 17-18, 24, L. verruculapeza sp. n. p. 83 pl. viii figs. 12-13, 15-16, 19-20 Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. gool. Surv. No. 334-C 1960.

Lloydiinae see Prolloydia gen, n.

Lonchocephalidae see Talbotinella gen. n.

Lonchocephalus sp. undet. p. 818 pl. xevii fig. 32, Lochman & Hu.

Lenchodomas megeheei Decker 1931 holotype refigured p. 215 pl. i figs. 1-4, pl. ii fig. 1, Sutherland & Amsden.

Lonchodomas pennatus (La Touche) [holotype figured] showing muscle scars p. 82 pl. xi figs. 2, 5, 8-12, Dean (1).

Lonchodomas portlocki (Barrande 1846) with synonyms Ampyx tetragomus Angelin, Raphiophorus depressus A. and A. t. var. gigas Linnarsson p. 169 pl. xxxiii fig. 8, pl. xxxv fig. 4, Kielan (1).

Lonchodomas rostratus (Sars et Boeck) var. taimyrica var. n. p. 31 (pl. vi figs. 6-7) Ordovician (Middle) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 (pls. in 14) 1960.

Lonchodomas sp. p. 30, Bogdanov & Chugaeva.

Lonchodomus sp. p. 39 pl. v fig. 2, Kobayashi (1).

Longiproetus 800 Proetus.

Lorenzella Kobayashi discussed, L. ? taitzuensis Endo excluded p. 388, L. lilia E. discussed p. 389; L. quadrata K. type of Cyclolorenzella gen. n. [q.v.] p. 389, Kobayashi (4).

Lotagnostus grandis Lermontova msc. [nom. nud.] pl. ii figs. 16-17, Chernysheva et al.

Lusatiops cf. morini n. sp. [nom. nud.] p. 81, Hupé (3).

Macrobole see Cyrtosymbole.

Maja Lomovitskaya 1955 replaced by Erdelia gen. n. p. 145, Chernysheva et al.

Maladidae [recte Maladiinae] Hupé a synonym of Monkaspidae p. 236, Kobayashi (3).

Maladiodella aff. splendens Endo p. 107, ÖPIK in Öpik & Pritchard.

Maladioidella may belong to Aphelaspidinao subfam. n. p. 80, Palmer (1).

Maladioides discussed and excluded from Elviniidae p. 64, Palmer (1).

Malykania Suvorova referred to Jakutidae (Malykaniae) subtam. n. p. 108, M. gribovae S. holotype figured p. 108 pl. ix figs. 9-15, pl. x figs. 1-9, text-fig. 36, M. grandis S. holotype figured p. 111 pl. x figs. 10-12 text-fig. 37, M. nochtujensie sp. n. with young stages p. 114 pl. x figs. 13-18, pl. xi figs. 1-19, text-figs. 38-39 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 34 1960.

Malykaninae subfam. n. [Jakutidae] p. 107 for Malykania and provisionally Mundocephalina, Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Manaspis gen. n. [Jakutidae] for type M. oblittera sp. n. p. 207 pl. i figs. 7-9 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Mansuyella Endo 1937 a synonym of Shirakiella p. 390, Kobayashi (4).

Mansuyia Sun 1924 discussed p. 363, M. hopeiensis sp. n. for M. orientalis Sun 1934 partim (pl. 3 figs. 7i-j) China p. 364 text-fig. 5a, M. trigonalis sp. n. S. Korea Cambrian (Upper) p. 364 pl. xx fig. 20 [sio], pl. xxi fig. 18, text-fig. 5c, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Mapania Resser & Endo discussed and referred tentatively to Liopariinae subfam. n. p. 392, Kobayashi (4),

Mareda mukazegata Kobayashi p. 407 pl. xix figs. 31-32 text-fig. 14b, Kobayashi (4).

Maryvillia sp. undet. p. 818 pl. xevii figs. 27-28, Lochman & Hu.

Mcnairia inornata Deiss p. 827 pl. c figs. 39-40, Lochman & Hu.

Megadunderbergia Kobayashi considered indeterminate [see also Dunderbergia] p. 66, Palmer (1).

Megalaspis see Megistaspis, M. centron (Leuchtenberg) discussed p. 185, M. latilimbata Angelin a nom. dub. p. 197, Bohlin.

Megalaspis gladiator, M. cf. g., M. sp. table 1 p. 6, Volk.

Megamansuyia glabra (Endo) listed p. 333, Kobayashi (4).

Megistaspidella see Megistaspis.

Megistaspis Jaanusson 1956 discussed p. 166 and spp. grouped into: A (1) M. lawrowi (Schmidt 1906) lectotype chosen p. 168 pl. iii figs. 2-4 text-figs. 10, ? 11; (2) M. heros (Dalman 1828) lectotype chosen and figured p. 170 pl. ii fig. 4, pl. iv figs. 2-3, pl. v fig. 1, pl. vi figs. 1-2, text-figs. 12-13, 16A, c, M. heros as pn. for M. heros of Schmidt 1898 and 1906 p. 174 pl. ii fig. 2, pl. v fig. 2, pl. vii figs. 1-8, text-figs. 15, 16B, p. E; B M. (Megistaspidella) including (3) Megalaspis rudis Angelin 1854 nom. dub., M. (M.) grandis (Sars 1835) lectotype chosen and figured p. 179 pl. iii fig. 1, M. (M.) acuticauda (A. 1854) lectotype chosen and figured [excluding Schmidt 1906] p. 181 pl. viii figs. 1-7 text-figs. 17-20, M. (M.) spinulata sp. n. for Megalaspis a. var. typica of S. 1906 p. 186; (4) M. (M.) convexa sp. n. for M. grandis var. lata and var. rudis of Törnquist 1884 p. 186 pl. ix figs. 1-5 text-figs. 21-22, M. (M.) sp. of Bohlin 1949 and 1955 p. 189 pl. ii fig. 3, pl. x figs. 1-5, text-figs. 23-24, M. bombifrons sp. n. p. 192 pl. ii figs. 1, pl. xi figs. 1-7 text-figs. 7, 25-26, M. (M.) ef. b. pl. i figs. 2-3, pl. iv fig. 1, M. (M.) obtusicauda (B. 1955) p. 197 text-figs. 6, ? 274, M. (M.) sigas (A. 1851) p. 198 pl. xi figs. 1-5 text-figs. 278, c. 28 Ordovician (Lower) Scandinavia and Baltic [see also Megalaspis], Bohlin Bull. geol. Instn Univ. Uppsala 38 2 1960.

Meneviella sp. p. 222 pl. xvii fig. 11, Repina.

Menocephalities Kobayashi discussed and referred to Menocephalitinae subfam. n. p. 381, Kobayashi (4),

Menocephalitinae subfam. n. [Solenopleuridae] p. 380 for Menocephalites, Paramenocephalites gen. n., Liaotungia, Eilura, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Menocephalus admeta Walcott type of Paramenocephalites gen. n. p. 381, Kobayashi (4).

Menomoniidae see Josina gen. n.

Menoparia referred to Proapatokephalidae fam. n. p. 33, Rozova (2).

Mesoparia ord. n. for Calymenidae and Homalonotidae p. 38, Lisbôa Rev. Escola de Minas, Ouro Preto 16 4 1951.

Metacalymene baylei (Barr. 1846)—Calymene tenera B. is apparently identical p. 114, Vaněk,

Metagraulos [Kobayashi]—Yabeia Resser & Endo probably a synonym p. 387, Kobayashi (4).

? Metaphillipsia seminiferus (Phillips) p. 311, Wilson.

cf. Metaphillipsia seminifera (Phillips) p. 129,

Metapilekia martellii (Kobayashi) for Metopolichas 7 m. K. 1934 p. 260, M. sp. nov. p. 261 pl. xiii fig. 26, Kobayashi (3).

Metascutellum gen. n. [Scutelluidae] p. 184 (259) for type M. pustulatum (Barrande 1846) for Bronteus pustulatus B. lectotype chosen and figd. [excluding records of B. 1852 partim pl. 48 figs. 13-16 and 1872 to "Scutellum" multiverrucatum sp. n.] with synonym S. gaudryi Prantl & Přibyl 1947 holotype figd. p. 185 pl. xix figs. 6-14 text-fig. 57 Devonian (Lower) Bohemia, Snajdr Rozpr. Úst. úst. geol. 26 1960.

Metashantungia 800 Shantungia.

Metisaspina Sivov may belong to Liopariinas subfam. n. p. 393, Kobayashi (4).

Metopolichas f martellii Kobayashi referred to Metapilekia p. 260, Kobayashi (3).

Micmacca Matthew discussed p. 64, M. (Micmacca) discussed p. 65, M. (M.) tumida sp. n. p. 66 pl. vii figs. 4-8 text-fig. 21 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Micmacca meridionalis n. sp. [nom. nud.] p. 81, Hupé (3).

Micragnostus see Geragnostus.

Micragnostus coreanicus sp. n. p. 234 pl. xiii figs. 7-11 Ordovician (Lower) Korea (South), Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Mic M. tet 1-3, ?

[19

Mis

and S

Mid for ty B. asj and f M. zij figd. [dormi

figs.
? M.
figs.
choses
(Lowe
1960.

Mit type Camb **Hu** J

Koba

Miconser um p (Nove 334–0

Mi Mo

Hu.

p. 23

Mo Malae branc

Mo

Mo aspis p. 26

classi Carin Mo

Mond kaspi Mc ŋ

i to

(4).

p.

kyo

eno.

. n.

uro

nera

ndo

311,

129,

chas

. 26,

) for

teus ding

1872

nym l. p.

wer)

iinae

d to

acca)

l. vii

.S.R.

Inst.

. 81,

figs.

yashi

Microparia speciosa Hawle & Corda 1847 with synonyms Aeglina ? oblongula Angelin type refigured and Symphysurus superstes Olin 1906 restr. [non pl. 3 fig. 6] of which lectotype selected and figured p. 88 pl. x figs. 4-9, Kielan (1).

Microphillipsia gen. n. [Phillipsidae] p. 5 for type M. tetraptera sp. n. p. 6 pl. i figs. l, 4-9, ? 2-3, text-figs. l-3, ? 4 Permian (Middle) Sicily, Ruggieri Mem. Inst. geol. Univ. Padova 21 1959.

Microscutellum gen. n. [Scutelluidae] p. 92 (252) for type Bronteus haulei Barrande 1852 with synonym B. asperulus Hawle & Corda [both lectotypes chosen and figd.] p. 93 pl. xxxiv figs. 8-13 text-fig. 31, M. zippei (B. 1846) for B. z. B. lectotype chosen and figd. [excluding B. 1852 pl. 45 figs. 22-23 to Paralejurus dormitzeri] p. 95 pl. xxviii figs. 8-10, pl. xxix fig. 3 [figs. include another "lectotype" of B. asperulus], 7 M. tenellum (B. 1852) for B. t. B. restr. (non pl. 47 figs. 36-38 to Paralejurus campanifer) lectotype chosen and figd. p. 97 pl. xxxv figs. 15-16 Devonian (Lower) Bohemia, Snajdr Rozpr. Úst. úst. geol. 26 1960.

Mimana euryrachis Kobayashi listed p. 333, Kobayashi (4).

Minicephalus gen. n. [Lecanopygidae] p. 816 for type M. primus sp. n. p. 817 pl. xcvii figs. 42-52 Cambrian (Upper) U.S.A. (Wyoming), Lochman & Hu J. Paleont. 34 5 1960.

Minupeltis gen. n. [Ptychoparioidea] for type M. conservator sp. n. with pathological paratype cranidium p. 98 pl. x figs. 11-12 Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Miomera accepted for Agnostina and Eodiscina p. 233, Kobayashi (3).

Miraspis mira (Barrande) text-fig. 615, Müller.

Modocia—relations to Ithyektyphus discussed, M. nevadensis Palmer referred to I. p. 829, Lochman & Hu.

Monkaspidae Kobayashi discussed with synonym Maladiidae Hupé and including Eurekiinae H. as a branch p. 236, Kobayashi (3).

Monkaspidae see Yosimuraspis gen. n.

Monkaspis Kobayashi discussed p. 237; Liaoningaspis and Kushanopyge [quae vide] may be synonyms p. 268-9, Kobayashi (3).

Monorakeidae Kramarenko 1952 discussed and classified into Monorakeinae and Elasmaspisinae & Carinopyginae subfamm. n. p. 6, Balashova (1).

Monorakeinae discussed and taken to include Monorakos, Evenkaspis, Parevenkaspis, Ceratevenkaspis, Trypaulites p. 7, Balashova (1).

Monorakeinae see Elasmaspis gen. n.

Monorakos magnus Kramarenko p. 8 [pl. iv fig. 5], M. mutabilis K. p. 9 [pl. iv fig. 6], M. ? nordicus sp. n. [no ? on pl. expl.] p. 10 [pl. iv figs. 3-4], M. zlobini sp. n. p. 11 [pl. iv figs. 1-26] Ordovician (Middle) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Moretaeva n. gen. [nom. nud.] [Strenusevinae] p. 77, Hupé (3).

Morosa gen. n. [Ptychoparioidea (? Housiidae)] p. 98 for type M. longispina p. 99 pl. x figs. 15-17 text-fig. 22 (reconstr.) Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334—C 1960.

Mundocephalina Repina referred provisionally to Malykaninae subfam. n. p. 107, Suvorova.

Myopsolenus see Protolenid.

Mystrocephala gen. n. [Brachymetopidae] p. 413 for type Cordania pulchra Cooper & Cloud 1938 holotype figured p. 414 pl. liii figs. 15-16, 19, 22-26, pl. liv figs. 1-5 text-fig. 1a, M. rara (Stumm 1953) [for C. r. S.] holotype figured p. 415 pl. lii figs. 10, 12, 15, M. varicella (Hall & Clarke 1888) lectotype selected and figured p. 415 pl. liv figs. 6-10, 12 Devonian (Lower and Middle), Whittington J. Paleont. 34 3 1960.

Namanoia Lermontova discussed, N. namanensis L. lectotype selected p. 153 pl. xiv figs. 1-9 text-fig. 50, N. evetasica sp. n. p. 156 pl. xiv figs. 10-15 textfigs. 51-52, N. chorincensis sp. n. p. 160 pl. xiv fig. 16 text-fig. 53 Cambrian (Lower) U.S.S.R. (Siberian Platform), N. tumida L. nom. nud. p. 153, Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Namanoidae discussed p. 152, Suvorova.

Namuropyge R. & E. Richter excluded from Brachymetopidae p. 408, Whittington (1).

Neoasaphus see Asaphus.

Neogriffithides ? siculus Gemm. for Phillipsia s. G. p. 3, Ruggieri.

Neopagetina gen. n. [Pagetiidae] (nom. n. for Pagetina Lermontova 1940) p. 56 for type P. rjons-nitzkii L. 1940 pl. ii figs. 20-21 Cambrian (Lower and Middle) U.S.S.R., Pokrovskaya Principles of Palaeontology. Arthropods—Trilobitomorphs and Crustaceans 1960.

Neoprobolium oklahomae (Richardson) for Dalmanites o. R. p. 53 figs. 1-2 age considered Devonian (Lower), Frederickson.

Neoproetus verrucosus Gemm. for Griffithides v. G. p. 4, Ruggieri.

Neoredlichiidae discussed with table of genera p. 21, Suvorova.

Neoredlichiidae see Tungusella gen. n.

Nericia discussed and may belong to Pterocephaliidae p. 80, N. septemdentata possibly to Pterocephaliinae p. 84, Palmer (1). Nescuretus Hioks 1872 discussed with synonym Synhomalonotus Pompeckj and provisionally retained in Calymenidae p. 138, N. grandior sp. n. for Calymene sp. Whittard 1931 p. 141 pl. xx figs. 1-2, N. parvifrons (Salter) holotype figured with synonym N. ? elongatus H. p. 142 pl. xix figs. 1-6, N. brevisulcus sp. n. p. 146 pl. xix figs. 7-14, pl. xx fig. 3, N. complanatus sp. n. possibly including O. pompeckji Kobayashi 1937 partim (pl. 6 fig. 24) p. 147 pl. xx figs. 4-5, N. murchisoni (S.) lectotype figured [with synonyms N. ramseyensis Hicks (lectotype selected and figured p. 139), N. recurvatus, N. ? e. obesus and probably N. quadratus H. partim (pl. 3 figs. 11, 13), ? C. p. K. 1937 partim (pl. 6 fig. 23), ? S. kobayashii Harrington & Leanza 1957] p. 148 pl. xx figs. 6-15, pl. xxi figs. 1-2 Ordovician (Lower), N. bullatus sp. n. p. 150 pl. xxiii figs. 9-11 Ordovician (Middle) Britain (Shropshire), Whittard Palaeontogr. Soc. [Monogr.]

Nevadella sp. p. 88 pl. iv fig. 3 Cambrian (Lower) Spitsbergen; N. eucharie (Walcott) and N. perfecta (W.) considered conspecific by R. V. Best p. 89, Kielan (2).

Nileus armadillo (Dalman) [small trilobite resembling] p. 275, Stanton.

Nileus sp. table 1 p. 6, Volk.

Niobe cf. emarginula Angelin p. 217, Tjernvik.

Niobe lata Angelin referred to Opsimasaphus gen. n. [q.v.] p. 78, Kielan (1).

Niobella bohlini Tjernvik p. 217, Tjernvik.

Nodiceps Suvorova discussed p. 123, N. onerosa S. holotype figured p. 124 pl. xii figs. 2-6 text-figs. 40-41, Suvorova.

Novaspis albida (Reed 1914) p. 175 pl. xxxiv fig. 3, pl. xxxvi figs. 1-5, text-fig. 50, N. sp. p. 178 pl. xxxv fig. 3, Kielan (1).

Octobronteus Weber 1945 discussed p. 56 (246), O. khodalevitchi W. 1945 lectotype chosen and figured p. 56 pl. xxxvi figs. 3-4 text-figs. 17A-B, Snaidr.

Odontopleura ovata text-fig. 4a, Whittington (3).

Odontopleura ? sp. Strnad 1956 p. 344, Strnad (1).

Odontepleurina referred to Lichida p. 233, Kobayashi (3).

Oedicybele kingi Whittington 1938 p. 157 pl. xxv fig. 4, Kielan (1).

Oedorhachis Resser 1938 discussed p. 12, O. australis sp. n. p. 13 pl. i fig. 8 Cambrian (Middle) Argentina (Mendoza); O. boltonensis, tennesseeensis, mesieri and greendalensis R. spp. may belong to Pseudagnostus, Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Ogmasaphus Jaanusson 1953 discussed and may be only a subgenus of Asaphus p. 235, O. jaanussoni sp. n. p. 237 pl. ix figs. 1-3, O. kiaeri sp. n. p. 239 pl. x figs. 1-12, O. multistriatus sp. n. p. 241 pl. vi figs. 13-17, O. stoermeri sp. n. p. 242 pl. xi figs. 1-6, O. sp. p. 244 pl. xii fig. 9 Ordovician (Middle) Norway (Oslo), Henningsmeen Norsk geol. Tidsskr. 40 3-4 1960.

Ogmocnemis gen. n. [Proetinae] p. 69 for type O. irregularis sp. n. Ordovician (Upper) Poland (Lysa Gora) p. 70 pl. iii figs. 5-9, pl. iv figs. 8-9, pl. xxvi fig. 1 text-fig. 17; Trilobites asellus Esmark, Forbesia brevifrons Angelin, Proetus modestus Tornquist, P. remotus Warburg, P. subornatus Cooper & Kindle referred to O. p. 69, Kielan Palaeont. polon. No. 11 1960.

Ogygia ? apiculata Linnarsson 1866 a synonym of Liocnemis recurvus (L.) p. 121, Kielan (1).

Ogygiocaris Angelin 1854 discussed and restricted to Scandinavian forms p. 213, stratigraphical range p. 214, glabellar segmentation p. 212 text-fig. 2, O. dilatata (Brünnich 1781) lectotype selected and figured with synonyms var. strems Angelin 1878 [lectotype selected and figured] and Asaphus debuchianus Brongniart 1817 p. 217 pl. ii figs. 1-7, pl. ii figs. 1-6 text-fig. 4 (reconstr.), O. striolata sp. 1. p. 221, D. c. striolata subsp. n. p. 222 pl. iii figs. 1-3, 11, O. sarsi A. 1878 discussed with notes on ontogeny p. 223, O. s. sarsi A. 1878 neotype [called lectotype on pl. expl.] selected and figured p. 225 pl. iv figs. 1-10 text-fig. 5 (reconstr.), O. s. regina subsp. n. [including O. d. sarsi partim of Hadding 1913 (text-fig. 23) & of Størmer 1953 (pl. 2) and O. lata partim of Thorslund 1935 (pl. 1 fig. 1)] p. 227 pl. v figs. 1-8, pl. vi fig. 5. o. s. delicata subsp. n. p. 228 pl. vi figs. 1-4, 6-12 Ordovician (Middle) Norway (Oslo), O. s. lata H. 1913 lectotype selected for O. d. l. H. [including O. d. sarsi of H. 1913 (non text-fig. 23) and O. d. ström of H. 1913 p. 229 pl. v fig. 9; O. araiorhachis Harrington & Leanza, O. ? plana and O. ? inflexicostata Thoral excluded from O. p. 214, Henningsmeen Norsk gool. Tidsskr. 40 3-4 1960.

Ogygopsis recorded from L. Cambrian of California and Nevada p. 2071, Nelson.

Ogygopsis klotzi (Rom.) in apparent Salter'sche Einbettung p. 104 text-figs. 1-2, Lehmann.

Olenaspella vertical range discussed, O. [2 new spp.] p. 44, Lochman-Balk.

Olenellidae discussed p. 17, Suvorova.

Olenellus svalbardensis sp. n. p. 84 pl. i figs. 1-3, pl. ii figs. 1-5, pl. iv figs. 1-2 Cambriaa (Lower) Spitsbergen, O. truemani Walcott discussed p. 88, Kielan Studia geol. polon. 4 1960.

Olenellus sp. text-fig. 4i, Whittington (3).

Olenina Poletaeva [not seen]—Arinichevia P. s synonym p. 26n, Ivshin.

Olenoidae see Prospatokephalidae fam. n.

1-3, p 5-7, 6 U.S.S S.S.S.

0. sp.

[19

Ole

Oler 18, 2 sp. n. U.S.A Surv.

Oligeott) 1 (1).

One distinct (B.) le O. grapl. xi: p. 133 p. 136 Storm verrue from (ficatio

Onn

579в,

Ops O. jac includ (pl. 31 O. late choser pl. viii 1906 No. 11

> Ord p. 371 hania

> > Orlo

Orde

Sivov Kobay Orm (Salter

figured pl. xvi l. H. n. p. l sp. n. 131 p [Monog

otar sp. St. figs. 1 Devon Přír. Č x

gs. O.

ay 3-4

nd

pl. rk,

å on.

of

ted

and 878 hus

-7, n. 10, 11,

eny

ype -10

ing

of and . 5, 3-12

913 d. i of ton

oral

rsk

mia

che

new

1.3,

rian

. 8

Olenoides convexus Lermontova p. 197 pl. ix figs. 1-3, pl. xviii fig. 4, O. entis sp. n. p. 198 pl. xviii figs. 5-7, O. sp. p. 198 pl. xvii fig. 12 Cambrian (Middle) U.S.S.R. (E. Sayan), Repina Regional Strat. 8.S.S.R. 4

of. Olenoides incertus Shaw p. 827 pl. c fig. 41, 0. sp. undet. p. 799, Lochman & Hu.

Olenus? wilsoni Henningsmoen p. 79 pl. vi figs. 18, 20-22 text-fig. 15 (reconstr.), O.? granulatus 59. n. p. 79 pl. vi figs. 23-27 Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334—C 1960.

Oligometopus Resser discussed, O. breviceps (Walcott) holotype figured p. 100 pl. x figs. 19-20, Palmer (1).

Onnia Bancroft 1933 discussed with table of distinctions from Broeggerolithus p. 127, O. ? cobboldi (B.) lectotype selected p. 128 pl. xix figs. 1, 3, 12, 0. gracilis (B.) lectotype selected and figured p. 130 pl. xix figs. 2, 7, O. superba (B.) cephalic ontogeny p. 133 pl. xix figs. 4-6, 8-9, 11, 13-14, O. aff. s. (B.) p. 136 pl. xix fig. 10; Cryptolithus cf. portlocki of Stermer 1930 probably an O. related to O. s., O. servicosa Rusconi 1956 should probably be excluded from O. p. 138, O. c. of G. & H. Termier 1950 identification suspect p. 137, Dean (1).

Onnia ornata (Sternberg) text-figs. 514 pars, 579B, Müller.

Opsimasaphus gen. n. [Asaphinae] p. 75 for type O. jaanussoni sp. n. Ordovician (Upper) Bornholm including Asaphus nobilis of Barrande 1852 partim (pl. 31 figs. 2-3, 6-8) p. 79 pl. vii figs. 1-2 text-fig. 21, O. latus (Angelin 1851) for Niobe lata A. lectotype chosen and figured p. 78 pl. vii figs. 1-2, pl. vii fig. 3, pl. viii fig. 4 text-fig. 20, O. sp. for A. ingens of Olin 1906 p. 81 pl. viii fig. 3, Kielan Palaeont. polon. No. 11 1960.

Ordosia Lu discussed p. 371, Kobayashi (4).

Ordosiinae discussed and referred to Pagodiidae p. 371; Taitzuia, Poshania, Lichengia and Peichiaehania referred, Kobayashi (4).

Orloviella elandiensis [Egorova] and O. conica Sivov may belong to Liopariinae subfam. n. p. 392, Kobayashi (4).

Ormathops Delo 1935 discussed, O. nicholsoni (Salter) for Phacops n. S. lectotype selected and figured with synonym O. barroisi (Klouček) p. 128 pl. xvi figs. 6-7, O. llanvirnensis (Hicks) for Phacops l. H. syntypes figured may be a synonym of O. n. p. 130 pl. xvi figs. 8-9, pl. xviii fig. 1, O. alatus sp. n. Ordovician (Middle) Britain (Shropshire) p. 131 pl. xvi fig. 10, Whittard Palaeontogr. Soc. [Monogr.] 113 (for 1959) 1960.

Otarion (O.) couvinianum sp. n. for Cyphaspides sp. Strnad 1956 [recte 1957] p. 340 (349) pl. iv figs. 14-17, pl. ii fig. 10 pars, 3 text-figs. [p. 341] Devonian (Middle) Czechoslovakia (Silesia), Strnad Přir. Čas. Slezský 21 3 1960. Otarion of. megalops (McCoy), O. sp. p. 177, Squirrell & Tucker.

Otarion aff. novella Barrande, O. sp. p. 156, Cavet.

"Otarion" tenuis sp. n. p. 63 pl. ii figs. 1-2 text-fig. 15, "O." sp. a p. 179 pl. viii fig. 6, O. sp. b p. 180 pl. ii figs. 8-9, "O." sp. c p. 181 pl. vii fig. 4, pl. viii fig. 5 Ordovician (Upper) Poland (Lysa Gora), Kielan Palaeont. polon. No. 11 1960.

Otarion (O.) n. sp. [undescr.], O. (Coignouina) davidsoni Barr. p. 350, Chlupáč & Vaněk.

Otarion sp. [compared with O. barrandei] p. 45,

Otarion sp. p. 350, Tomczykowa.

Otarion sp. p. 412, Whittington (1).

Ouijjania referred to Bigotinopsidae fam. n. p. 77, Hupé (3).

Pagetiellus lenaicus (Toll) from Morocco p. 77, Hupė (3).

Pagetiellus sibiricus sp. n. p. 175 pl. xiii figs. 11-14 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Pagetiidae see Neopagetina gen. n.

Pagetina Lermontova 1940 renamed Neopagetina gen. n. p. 56, Chernysheva et al.

Pagetina primaeva Lerm. pl. v fig. 14 pars, Suvorova.

Pagodia Walcott 1905 discussed with list of spp. p. 366, P. coreanica sp. n. p. 367 pl. xix figs. 27-29 text-fig. 60 Cambrian (Upper) S. Korea, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Pagodiidae taken to include Pagodiinae and Ordosiinae p. 366, Kobayashi (4).

Pagodiinae—Chuangiopsis referred p. 370, Kobayashi (4).

Paladin Weller 1936 discussed p. 34, P. eichwaldi (Fischer 1825) p. 35 pl. vii fig. 1 pars A, pl. viii fig. 1 pars A, pl. viii fig. 1 pars A, pl. ix figs. 1-4, pl. x figs. 1-2, P. mucronatus (McCoy 1844)—Pseudophillipsia croatica Schréter 1948 a synonym p. 47n, Bouček & Přibyl.

Palaeolenella gen n. [Palaeolenidae] p. 214 for type P. artifeza sp. n. p. 215 pl. x figs. 1-3, P. doliara sp. n. p. 215 pl. x figs. 4-5 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.R. 4 1960.

Palaeolenidae see Palaeolenella gen. n.

Palaeophillipsia see Palaeophyllipsia.

Palaeophyllipsia [so spelt at descr. and pl. expl., elsewhere Palaeophillipsia] gen. n. [compared with Phillipsia], P. japonica sp. n. p. 26 pl. i fig. 1 text-fig. (p. 28), P. i kitakamiensis sp. n. p. 29 pl. i figs. 2a-o Devonian (Upper) Japan (Iwate), Sugiyama & Okano Res. Bull. [Stud.] geol. miner. Inst. Tokyo Bunrika Daigaku No. 1 1944.

Palpebralia referred to Drevermanniinae subfam. n. p. 137, Chernysheva et al.

Panderia megalophthalma Linnarsson 1869 including Symphysurus superstes Olin 1906 partim (pl. 3 fig. 6) p. 93 pl. xii fig. 6, Kielan (1).

Paraaulacopleura see Aulacopleura.

of. Parabasilicus powisi (Murchison), ? P. d. p. 167, Dean (3).

Parabolinoides expansus Nelson p. 811 pl. xev figs. 32-35, P. cf. P. e. p. 798, Lochman & Hu.

Paraceratopyge Poletayeva 1945 [not seen] [Ceratopyginae] discussed p. 104, P. kusnezkiensis P. text-figs. 193a-6, Chernysheva et al.

Paracolpocoryphe dubia n. gen. n. sp. p. 1537 [nom. nud.—near to Colpocoryphe grandis Snajdr. p. 1539], Destombes et al.

Paracybeloides cf. loveni (Linnarsson) p. 334,

Paradoxides bohemicus text-fig. 4f, Whittington (3).

Paradoxides gracilis (Boeck) text-fig. 532B, Müller.

Paragraulos Lu 1941 discussed with synonym Inouyops Resser p. 385, Kobayashi (4).

Parahousia gen. n. [Housiidae] p. 76 for type P. constricts sp. n. p. 77 pl. vii figs. 16-18 text-fig. 13 (reconstr.) Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Parairvingella see Irvingella.

Parairvingella hamburgensis Resser referred to Elvinia [quod vide] p. 71, Palmer (1).

Parakingaspidoides n. gen. [nom. nud.] [Palaeo-lenidae] p. 77, P. truncatus n. sp. [nom. nud.] p. 81,

Parakoldinia gen. n. [Plethopeltidae] p. 49 for type P. striata sp. n. p. 52 pl. vi figs. 1-6, P. plana sp. n. p. 54 pl. vii figs. 1-5, P. hemisphaerica sp. n. p. 55 pl. vii figs. 6-10, P. longa sp. n. p. 57 pl. vi figs. 13-18 Cambrian (Upper) U.S.S.R. (Salair), P. salairica (Sivov) p. 58 pl. vi figs. 7-12; P. gibbosa (Lermontova), P. globosa (Sivov), P. paichinensiformis (S.) transerred from Kingstonia p. 52, Rozova Trud. Inst. eol. geophiz. AN S.S.S.R. (Sib. Otd.) 5 1960.

Paralejurus Hawle & Corda 1847 discussed p. 187 (260), P. campanifer (Beyrich 1845) neotype chosen and figd. [including Bronteus tenellus of Barrande 1852 partim pl. 47 figs. 36-38] p. 189 pl. xxiii figs. 2-14, pl. xxiv figs. 1-2, pl. xxxvi fig. 8, text-figs. 58-59, P. brongniarti b. (B. 1846) with synonym B. pulcher Hawle & Corda 1847 both lectotypes chosen and figd. p. 193 pl. xxi figs. 9-14, pl. xxii figs. 1-13, pl. xxiii fig. 1, text-fig. 60, P. b. méhanensis subsp. n. p. 196 (260) pl. xxv figs. 10-11, P. dormitzeri d. (B. 1852) with synonym B. aulicus H. & C. 1847 both lectotypes chosen and figd. and including B. zippei of B. 1852 partim (pl. 45 figs. 22-23) but excluding B. d. of Le Maitre 1934 p. 196 pl. xxiv figs. 3-16, pl. xxv figs. 1-3, text-fig. 61, P. bohemicus Sp. n. p. 200 (261) pl. xxv figs. 4-5, P. richteri sp. n. p. 201 (261) pl. xxv figs. 6-9 Devonian (Lower) Bohemia, Snajdr Rozpr. Ust. úst. geol. 26 1960.

Paramansuyella glabra Endo type of Taipaikia gen, n. p. 398, Kobayashi (4).

Paramansuyella aff. puteata Endo p. 107, Örik in Öpik & Pritchard.

Paramenocephalites gen. n. [Menocephalitinae] p. 381 for type Menocephalus admeta Walcott 1995 text-fig. If Cambrian (Middle) China; Damesella nitida Rosser & Endo also referred to P., Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Paramenomonia Chu 1959 discussed and referred to Damesellidae (Paramenomoniinae) subfam. n. p. 354, P. conica C. text-fig. 2h, Kobayashi (4).

Paramenomoniinae subfam. n. [Damesellidae] for Paramenomonia Chu 1959 p. 354, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Paramicmacca Lermontova with synonym Acanthomicmacca Hupé, P. siberica L. lectotype chosen p. 68 pl. vii figs. 1-3 text-fig. 22, P. petropaulouskii sp. n. p. 72 pl. vii figs. 9-10 text-fig. 23, P. insolens sp. n. p. 73 pl. vii figs. 11-13 text-fig. 24 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 84 1960.

Parapoliella Chernysheva referred to Jakutinae p. 86, Suvorova.

Parapoliella obrutchevi (Lermontova) p. 206 pl. vi fig. 7, P. sulcata Chernysheva [nondescr.] p. 206 pl. vi figs. 1-6, pl. viii figs. 1-3, Repina.

Parashantungia see Shantungia.

Parasolenopleura Poletayeva [non Westergård] discussed p. 377, Kobayashi (4).

Pedinocephalina gen. n. [compared with Pedinocephalus and Coenaspis] p. 201 for type P. bella sp. n. p. 203 pl. i figs. 1-2, P. composita sp. n. p. 205 pl. i figs. 3-4 Cambrian (Upper) U.S.S.R. (Salair), Rozova Trud. Inst. geol. geophiz. AN S.S.S.R. (Sib. Otd.) 1 1960.

Pedinocephalina Rozova discussed and referred to Alokistocarinae p. 70, P. bella R. holotype figd. p. 72 pl. iii figs. 13-14, P. composita R. holotype figd. p. 74 pl. iii figs. 15, 15a, Rozova (2).

Pedinocephalus see Pedinocephalina gen. n.

Peichiashania Chang discussed and referred to Ordosiinae p. 372, Kobayashi (4).

Permoproetus postcarbonarius Gemm. for Proetus p. G. p. 3, Ruggieri.

Peronopsinae see Stigmagnostus gen. n.

Peronopsis ultima sp. n. p. 13 pl. i fig. 13 Cambrian (Middle) Argentina (Mendoza) [may include cephala referred to Baltagnostus hospitus sp. n. q.v.], Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Pesaia avia Sivov referred to Pesaina p. 7, Ivshin.

Pesaina Poletaeva [not seen]—Pesaia avia Sivov referred p. 7, Ivshin.

Phace tubbles

Phace

Phace

Phace

Phace P. sp. r (Altai), trilobite

Phace 1931 p.

Phace Phace p. 1538

Phace chosen 3, 7, platernber

pl. iii fi

n. p. 8
oculeus
P. poste
(Upper
text-fig
figs. 6lectoty)
fig. 12,
8, P. s
simova
Rudny

Phae Hall, F

P. fugi Phac turco I P. (P.)

Phae Vaněk.

Phace (P.) as R. & H. p. 177, sp. p.

Phac

Phace [unillust Inst. T ia

K

p.

6.0

d]

05

ib.

to

to

tus

in.

TOV

Phacopid—Homalonotus barratti may be p. 104, stubblefield.

Phacopid glabella p. 270, Anderson.

Phacopid (undescribed) p. 892, Tillman.

Phacopidella liopyga Reinh. Richter text-fig. 515B,

Phacopidella (Denckmannites) micromma (A. Roemer 1852) p. 346, Strnad (1).

Phacopidella primitiva sp. n. p. 93 pl. ii figs. 5-11, P. sp. p. 96 pl. v fig. 13 Devonian (Middle) U.S.S.R. (Altai), Makrimova Devonian and Carboniferous trilobites of the Rudny Altai 1960.

Phacopidella (Denckmannites) thuringica Kegel 1931 p. 114 pl. xi figs. 5-9, Alberti.

Phacopidina apiculata (M'Coy) p. 168, Dean (3).

Phacopidina (Dreyfussina) cf. exophtalma Dreyfuss p. 1538, Destombes et al.

Phacopoid see Trilobite pygidium.

Phacops altaicus Tschernyschew 1893 lectotype chosen and figured p. 75 pl. i figs. 11-16, pl. iii figs. 3, 7, pl. v figs. 1-4, P. cf. a. p. 90 pl. iii figs. 8, P. sternbergi Hawle & Corda lectotype chosen p. 79 pl. iii fig. 2, pl. v figs. 5-10, P. potieri ulbensis subsp. a. p. 81 pl. ii figs. 12-18 Devonian (Middle), P. coulcus sp. n. p. 83 pl. iii fig. 4 Devonian (Middle), P. postaltaicus sp. n. p. 84 pl. viii figs. 9-13 Devonian (Upper), P. loganops sp. n. p. 86 pl. vii figs. 12-13, text-fig. 4 Devonian (Middle), P. aff. l. p. 91 pl. iv figs. 6-7, P. cristata var. pipa Hall & Clarke 1888 lectotype chosen p. 89, P. cf. c. var. p. p. 89 pl. iii fig. 12, P. aff. pronini Maksimova p. 92 pl. viii fig. 8, P. sp. p. 92 pl. iii fig. 13 U.S.S.R. (Altai), Maksimova Devonian and Carboniferous trilobites of the Rudny Altai 1960.

Phacops aff. batrachensis Whidb., P. aff. cristata Hall, P. ex gr. fecundus Barr., P. sp. p.121, Bel'akaya.

Phacops fecundus Barr., P. f. var. lyauteyi Lecointre, P. fugitivus B. p. 46, Cogney.

Phacops (P.) fecundus degener Barrande, P. (P.) turco R. & E. Richter, P. (P.) schlotheimi Bronn, P. (P.) aff. angusticeps Giebel p. 160, Cavet.

Phacops fecundus major Barr. p. 350, Chlupáč & Vapěk.

Phacops (P.) granulatus (Münster) p. 154 ff., P. (P.) accipitrinus a. (Phillips), P. (P.) wedekindi R. & E. Richter, P. (Cryphops?) ensae R. & E. R. p. 177, P. (C.?) wocklumeriae R. & E. R., P. (Dianops) sp. p. 178 pl. xxix fig. 4, **Selwood**.

Phacops nicholsoni Salter and P. llanvirnensis Hicks referred to Ormathops [q.v.] p. 128, 130, Whittard.

Phacops okanoti sp. n. Devonian Japan p. 24 [unillustr.], Sugiyama Res. Bull. [Stud.] geol. miner. Inst. Tokyo Bunrika Daigaku No. 1 1944.

VOL 97

Phacops okanoi Sugiyama p. 44, Okubo (2) [omitted Z. R. 93 11 (for 1956)].

Phacops quadrilineatus Angelin p. 21 [pl. v fig. 19], Balashova (1).

Phacops recurvus Linnarsson type of Liocnemis gen. n. p. 120, Kielan (1).

Phacops (P. ?) roemeri Geinitz 1853 p. 111 pl. xi fig. 4, P. (Eccryphope) cyclophthalmus (Walther 1907) p. 112 pl. xi figs. 1-3, P. (Plagiolaria) plagiophthalmus (Reinh. Richter 1865) p. 113, Alberti.

Phacops stokesi (Milne Edwards), P. sp. p. 177, Squirrell & Tucker.

Phacops truncatocaudatus Portlock type of Pterygometopus (Truncatometopus) subgen. n. p. 9, Pillet (3).

Phacops sp. indet. p. 346 pl. iv figs. 19-22, Strnad (1).

Phaëtonellus sp. (non planicaudus Barr.) p. 350, Chlupáč & Vaněk,

Phalacroma ? sp. p. 63 pl. iv fig. 13, Palmer (1).

Pharostoma Hawle & Corda 1847 discussed and referred to Calymenidae p. 132, P. pulchrum (Beyrich) so attributed p. 134 pl. xviii figs. 2-8, discussion of several spp. of P. p. 137, Whittard.

Pharostomidae see Ptychometopinae subfam. n.

Phillibole (Phillibole) aprathensis R. et E. Richter p. 101 pl. figs. 18-20, Strnad (2).

Phillibole (P.) coddonensis (H. Woodward) [of which P. (P.) culmica R. & E. Richter may be a synonym p. 273], P. (P.) aff. c. p. 272, P. (Liobole) glabra g. (Holzapfel) p. 271 pl. xii fig. 1, ? P. (L.) sp., P. ? (Cystispina) spatulata (H. W.) p. 271 pl. xii fig. 9, P. (? C.) sp. nov. p. 272 pl. xii fig. 2; Phillipsia [van-der-grachtii and polleni] H. W. 1894 have characteristics of P. s.s. p. 278, Prentice.

Phillipsia Portlock 1843 emend. Stubblefield 1952 discussed p. 23, Gruppe der Art P. gemmulifera recognised including also i.a. P. truncatula, P. atlaica, P. margaritifera, P. konincki, P. scabra, P. collinsi p. 25; P. margaritifera Roemer 1870 with possible synonym var. conferta Schwarzbach p. 25 pl. ii figs. 1-5, pl. iii figs. 1-2, pl. iv fig. 1, pl. v figs. 1-2, pl. vii fig. 1 pars B, P. collinsi Mitchell 1918 lectotype selected p. 28n, Bouček & Přibyl.

Phillipsia see Palaeophyllipsia gen. n.

Phillipsia see Phyllipsia.

Phillipsia altaica Weber 1937 p. 55 pl. ix figs. 9-11, P. buchtarmensis sp. n, including P. a. W. 1937 partim (pl. 5 figs. 6, 7) p. 57 pl. ix figs. 6, P. ? abscisa W. 1937 lectotype chosen p. 60 pl. ix fig. 1, P. aff. scabra Woodward 1883 p. 61 pl. ix fig. 12, P. ? minuta sp. n, for P. No. 18 of W. 1937 p. 62 pl. ix fig. 3, P. sp. p. 63 pl. ix fig. 4 Carboniferous (Lower) U.S.S.R. (Altai), Maksimova Devonian and Carboniferous trilobites of the Rudny Altai 1960.

Phillipsia ohmorensis sp. n. p. 133 pl. i figs. 1a-d, 2-4 Devonian (Upper) or Carboniferous (Lower) Japan (Iwate), Okubo Earth Science [Chikyū Kagaku] No. 4 1951.

Phillipsia sicula Gemmellaro referred tentatively to Neogrifithides p. 3, Ruggieri.

Phillipsia [van-der-grachtii and polleni] H. Woodward 1894 have characteristics of Phillibole s.s. p. 278, Prentice.

Phillipsia sp. ind. p. 183 text-fig. 2, Milolević.

" Phillipsia" sp. p. 4 pl. iii fig. 15, Armstrong.

Phillipsiid p. 309, Phillipsiid pygidium p. 310, Wilson.

Phillipsiidae see Microphillipsia gen. n.

Phillipsiinae discussed p. 22, Bouček & Přibyl.

Phillipsinella parabola (Barrande 1846) p. 72 pl. iv figs. 1-7, pl. v figs. 1-3 text-figs. 18-19; record of P. p. by Olin 1906 (p. 78) from Chasmops Beds belongs to new sp. of P. p. 38, **Kielan** (1).

Phillipsinellidae discussed and affinities may be with Proetacea pp. 47-48, Kielan (1).

Phoidagnostus solitariensis (Rusconi 1950) for Gallagnostus s. R. [re-interpreted as a pygidium] p. 14 pl. i fig. 15, Poulsen (2).

Phyllaspis Reinh. Richter 1863 nov. emend. discussed and referred tentatively to Tropidocoryphinae p. 119, P. raniceps R. R. 1863 nov. emend. [with synonym Proetus expansus R. R. 1865] neotype selected and figured p. 123 pl. xii figs. 1-6, text-figs. 3-5, Alberti.

Phyllipsia (pygidia) p. 49 (51), Stevanović.

Pinctus ? spp. p. 101 pl. x figs. 18, 21, Palmer (1).

Piriproetus see Proetus.

Placoparia (Hawleia) prantli sp. n. p. 125 pl. xxiii figs. 4-6 text-fig. 34 Ordovician (Upper) Poland (Lysa Gora), Kielan Palaeont. polon. No. 11 1960.

Placoparia cf. zippei Boeck p. 1538, Destombes et al.

Plagiolaria see Phacops.

Planaspis gen. n. [Aldoniidae] for type P. gelasinica sp. n. p. 221 pl. x fig. 9 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Planiscutellum R. & E. Richter 1956 discussed p. 57 (246), P. planum (Hawle & Corda 1847) lectotype chosen and figured p. 59 pl. i figs. 1-10, pl. ii figs. 1-2, text-figs. 18-20, Snajdr.

Plasiaspis Prantl & Přibyl 1948 diagnosis p. 122, considered to form with Dindymene and Cornovica gen. n. a compact group of Encrinuridae p. 121, Whittard.

Plasiaspis excluded from Dindymeninae p. 144, P. bohemica (Barrande) ventral cephalic sutures (separating a rostral plate) described p. 54 text-fig. 10, Kielan (1).

Plataspella Wilson discussed and considered separate from Iddingsia p. 96; I. similis of Bell et al. 122 partim (pl. 31 fig. 2) referred to P. p. 96, Palmer (1).

Platilichas [sie] cf. laxatus (McCoy) p. 334, Cave.

Platycalymene Shirley 1936 accepted as a genus p. 151, P. tasgarensis S. with muscle sears p. 152 pl. xxi figs. 3-12, P. duplicata (Murchison) [locality and horizon of holotype discussed] forma femina and forma mas both found in relief p. 154 pl. xxi figs. 13-15, P. d. var. parallela var. n. p. 157 pl. xxi fig. 16, P. d. of. p. p. 157 pl. xxi fig. 17 Ordovician (Middle) Britain (Shropshire), Whittard Palaeontogr. Soc. [Monogr.] 113 (for 1959) 1960.

Platylichas see Platilichas

Platysaukia gen. n. [Saukiinae] for type P. euryrachis sp. n. p. 407 pl. xix fig. 12, text-fig. 13 b Cambrian (Upper) S. Kores, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Platyscutellum Šnajdr 1958 discussed p. 118 (254), P. formosum f. (Barrande 1846) lectotype figd. [excluding records of R. & E. Richter 1943 and H. & G. Termier 1950 and with synonym Brontew berkeleyanus Hawle & Cords lectotype chosen and figd.] p. 119 pl. xvi figs. 1-4, text-fig. 41, P. aff. f. f. including B. planus of B. 1852 partim (pl. 48 fig. 1) pl. xv figs. 8-9, P. f. slivenecense subsp. n. p. 122 (254) pl. xv figs. 5-7, P. kutorgai (B. 1852) holotype figd. p. 123 pl. xiv figs. 6-8, pl. xv figs. 1-2, pl. xxxvi fig. 10, P. lochkovense sp. n. p. 124 (255) pl. xv figs. 3-4 text-fig. 42, P. viator (B. 1852) lectotype chosen and figd. [excluding B. 1852 pl. 47 figs. 21-22 to Breviscutellum inerme, Novák 1884 pl. 12 figs. 3-9, H. & G. Termier 1950, cf. Erben 1952] p. 126 pl. xvii figs. 2-8 text-fig. 43, P. cf. v. pl. xvii fig. 9, P. spiniferum (B. 1852) lectotype chosen and figd. with synonym B. sosia B. 1872 holotype figd. p. 129 ll. xvii figs. 5-8, pl. xvii fig. 1, text-fig. 44, P. magum (B. 1872) holotype figd. p. 131 pl. xviii fig. 1 Devonian (Lower) Bohemia, Snajdr Rozpr. Úst. úst. geol. 26 1960.

Plesiomegalaspis planilimbata (Angelin) pp. 216, 217, Tjernvik.

Plethometopus modestus p. 43, Lochman-Balk.

Plethopeltidae see Parakoldinia gen. n.

Plethopeltidae see Tolstotchichaspis gen. n.

Pliomerid gen. et sp. indt. p. 263 pl. xiii figs. 33-34, Kobayaxhi (3).

Pliomerops actinurus (Dalman) p. 217, Tjernvik.

Pliomerops sp. I, P. sp. II p. 30, Bogdanov & Chugaeva.

37 p (W. K Roso (Sib.

Caml

[18

Po

for t

Poi Repir

Por P. por figd. 45, F gervilli pl. xx p. 138 figd. (B. 18

Por allied Por kephs

Pos

P. lin

Prealata ( (recon 11, 14 ventre aff. A; P., Pi 1960.

[see al

Pric proetus Tomez

of. I

Apatok gen. phalus, Apatok Trud. 5 1960

77 7 7

44, res 10.

red

ner

152

ity ina

XXI

ian ogr.

3 b Sci.

54), igd. and

teus and . in-. 1; 254) figd.

**cxv**i

XV

ype 1-22

figs. 126

g. 9, figd.

129

gum

nian

216,

3-34,

nvik.

W &

Poletaevia gen. n. [† Proapatokephalidae] p. 36 for type P. shorica [schorica on pl. expl.] sp. n. p. 37 pl. v figs. 15-19 Ordovician (Lower) U.S.S.R. (W. Siberia); Pseudokainella † macarenae Harrington & Kay referred tentatively to Poletaevia p. 39, Rozova Trud. Inst. geol. geophiz. AN S.S.S.R. (Sib. Otd.) 5 1960.

Poliellaspis rotundata sp. n. p. 205 pl. x figs. 6-8 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Poliellina sp. p. 206 pl. xii figs. 4-5, pl. vii fig. 7, Repina.

Poroscutellum Snajdr 1958 discussed p. 132 (255), P. porosum (Barrande 1846) lectotype chosen and figd. p. 133 pl. xx figs. 1-11, pl. xxi fig. 1, text-fig. 45, P. infaustum (B. 1852) with synonym Bronteus gervilleicans B. 1872 (both holotypes figd.) p. 136 pl. xxi figs. 2-5, P. expectane (B. 1872) holotype figd. p. 138 pl. xxi figs. 6-8, P. indocile (B. 1872) holotype figd. p. 139 pl. xxxii figs. 7-9, P. acupunctatum (B. 1872) holotype figd. p. 139 pl. xix figs. 3-5, Snajdr.

Portentosus Egorova discussed and considered allied to Hukasawaia p. 243, Kobayashi (3).

Portentosus discussed and referred to Proapatokephalidae fam. n. p. 43, P. brevis Egorova p. 44 pl.iv figs. 9-14, Rozova (2).

Poshania Chang discussed and referred to Ordosiinae p. 372, Kobayashi (4).

Praedechenella Maksimova 1953 [not seen] p. 135, P. liniclivosa M. text-figs. 311a-6, Chernysheva et al.

Prantlia longula (Hawle & Corda 1847) p. 209, Přibyl.

Prehousia gen. n. [Housiidae] p. 77 for type P. alata sp. n. p. 78 pl. vii figs. 10, 12-13 text-fig. 14 (reconstr.), P. semicircularis sp. n. p. 78 pl. vii figs. 11, 14-15, 19 Cambrian (Upper) U.S.A. (Nevada), ventral cephalic sutures of P. text-fig. 8 p; N. gen. aff. Aphelaspis of Palmer & Hazzard 1956 belongs to P., Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Pricyclopyge prisca (Barrande) text-fig. 558 B [see also Cyclopyge], Müller.

Prionopeltis archiaci see Hypostom F.

Prionopeltis sp. p. 350 [altered by hand to Cyphoproctus rugosus in offprint from author p. 352], Tomczykowa.

cf. Proagnostus sp. undet. p. 823, Lochman & Hu.

Prospatokephalidae fam. n. [Olenoidae] p. 33 for Apatokephalus, Proapatokephalus, Proapatokephalus, Diplapatokephalus, Artokephalus, Apatokephalina, Portentosus, Apatokephaloides, † Poletaevia gen. n. p. 33, Rozova Trud. Inst. geol. geophiz. AN S.S.S.R. (Sib. Otd.) 5 1960.

Proapatokephalops gen. n. [Proapatokephalidae] p. 39 for type P. oblongus sp. n. p. 41 pl. iv figs. 1-6, P. latus sp. n. p. 42 pl. iv figs. 7-8 Cambrian (Upper) U.S.S.R. (Salair), Rozova Trud. Inst. geol. geophiz. AN S.S.S.R. (Sib. Otd.) 5 1960.

Proapatokephalus referred to Proapatokephalidae fam. n. p. 33, Rozova (2),

Proasaphiscus sp. p. 209 pl. ix fig. 8, Repina.

Probolium ? sp. α p. 97 pl. iv fig. 5, pl. vii fig. 14, P. ? sp. β p. 98 pl. iv [sic] figs. 1-3 Devonian (Middle) U.S.S.R. (Altai), Maksimova,

Proceratocephala terribilis bituberculata subsp. n. p. 107 pl. iii fig. 3, pl. xvi fig. 1 Ordovician (Upper) Poland (Lysa Gora), Kielan Palacont. polon. No. 11 1960.

Prochuangia Kobayashi 1935 discussed p. 361, Kobayashi (4).

Prodamesella Chang 1959 discussed p. 352, Kobayashi (4).

Proerbia Lermontova discussed p. 147, P. semota sp. n. p. 148 pl. xiii fig. 13 text-fig. 48 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 34 1960.

Proerbia torgaschinica sp. n. p. 203 pl. viii figs. 7-8, P. cf. prisca Lermontova pl. 203 pl. xv figs. 5-6, P. sp. p. 203 pl. viii fig. 3, P. sp. p. 204 pl. xiii fig. 8 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Proerbidae Hupé a synonym of Dinesidae Lermontova p. 139, Suvorova.

Proetacea—Phillipsinellidae may have affinities p. 48, Kielan (1).

Proetid p. 412, Whittington (1).

Proetid hypostomes figured by Barrande 1852 pls. 15-17 discussed and assigned p. 196, Přibyl.

Proetidella cf. fearnsidesi Bancroft p. 167, Dean (3).

Proetina Přibyl 1946 discussed with diagnosis and referred to Proetinae p. 209, P. latimargo P. 1946 including Proetius lusor of Barrande 1872 (non 1852) p. 210, Přibyl.

Proetinae-Proetina referred p. 210, Přibyl.

Proetinae see Ogmocnemis gen. n.

Proctus bohemicus Cord., P. natator Barr. p. 44, Cogney.

Proetus (Longiproetus) chamaeleo R. & E. Richter, P. (L.) cf. tenuimargo R. & E. R. p. 160, Cavet,

Proetus conspersus (Ang.) p. 350, Tomezykowa.

Proctus (Piriproctus) dormitans Reinh. Richter 1863 p. 117 pl. xii figs. 10-12 text-fig. 1, P. (Piriproctus) schaderthalensis sp. n. p. 118 text-fig. 2 Devonian (Middle) Germany (Thüringia); P. expansus R. R. a synonym of Phyllaspis runiceps R. R. p. 123, Alberti Mitt. geol. Staatsinst. Hamb. 29 1960.

Proctus latifrons McCoy, P. sp. p. 178, Squirrell & Tucker.

Proctus lazutkini N. Tchern., P. sp. p. 121, Bel'skaya.

Proetus modestus Törnquist, P. remotus Warburg, P. subornatus Cooper & Rindle referred to Ogmocnemis gen. n. p. 69, Kielan (1).

Proetus (P.) mořinensis sp. n. Silurian (Middle) Bohemia p. 178 (204) pl. ii figs. 1-5, P. (P.) retroflezus Barrande 1852 p. 206 pl. i figs. 11-16, pl. iii figs. 1-2, pl. iv fig. 11, P. (P.) signatus Lindström 1885 lectotype selected p. 182 (207) pl. i figs. 17-22, P. (subgen. ?) lusor B. 1852 p. 208, P. (Eremiproetus) insons B. 1852, P. (subgen. ?) moestus B. 1852 p. 209, P. (P.) ryckholti B. pl. ii fig. 6; P. l. of B. 1872 referred to Proetina latimargo p. 210; P. (Cornuproetus) duffesnoyi see Hypostom A, P. (C.) sp. see Hypostom B, P. (C.) complanatus B. see Hypostom C, P. (P.) neglectus B. [lectotype chosen p. 193] see Hypostom D, P. sp. see Hypostom E, Přibyl Sborn. Úst. úst. geol. 25 [Pal.] (for 1958) 1960.

Proetus (P.) oehlerti Bayle 1878 p. 39 pl. i figs. 6-7, P. (P.) sibiricus Tschernyschew 1893 lectotype chosen and figd. p. 40 pl. i figs. 9-10, P. (P.) bohemicus supraconvexus subsp. n. p. 41 pl. vi figs. 1-4 text-fig. 2, P. (P.) ex gr. bohemicus Hawle & Corda 1847 p. 44 pl. vii fig. 15, P. (P.) aff. retroflexus Barrande 1852 p. 45 pl. vi fig. 9, P. [P. (P.) on table 1] pseudocarbonicus sp. n. p. 46 pl. iii figs. 10-11, P. sp. for Dalmanites sp. T. 1893 p. 47 Devonian (Middle), P. (Semiproetus ?) aff. saryaensis Weber 1937 p. 48 pl. ix fig. 2, P. (S. ?) ussuilensis var. altaica W. 1937 p. 49 Carboniferous (Lower) U.S.S.R. (Altai); P. sp. T. 1893 referred to Eremiproetus cf. eremita p. 53, Maksimova Devonian and Carboniferous trilobites of the Rudny Altai 1960.

Proetus (P.) orbitatus Barr., P. (P.) myops B., P. (Cornuproetus) complanatus B., P. (C.) buchi (H. & C.), P. (Unguliproetus) unguloides B., P. (Eremiproetus) sp. p. 350, Chlupáč & Vaněk.

Proetus postcarbonarius Gemmellaro referred to Permoproetus p. 3, Ruggieri.

Proetus sp. pl. v figs. 18-19, Wilson & Majewske.

"Proetus" sp. a p. 181 pl. iii figs. 1-2, pl. xxi fig. 1, "P." sp. b p. 182 pl. xxi fig. 2, "P." sp. c p. 182 pl. xxv fig. 5 Ordovician (Upper) Poland (Lysa Gora), Kielan (1).

Proetus sp. indet. p. 269, Churkin & Langenheim,

Prohedinella erbiensis Sivov p. 213 pl. ix fig. 9, Repina.

Prolloydia gen. n. [Lloydiinae] for type P. orientalis sp. n. p. 365 pl. xx figs. 21-25, text-figs. 3 g-h Cambrian (Upper) S. Korea, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960. Promegalaspides kazachstanensis Balashova [nom. nud.] pl. v figs. 4-5, Chernysheva et al.

Prometeoraspis gen. n. [Tricrepiesphalidae] for type P. canotensis sp. n. p. 17 pl. i figs. 16-17 Cambrian (Middle) Argentina (Mendoza), Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Prosaukia Ulrich & Resser discussed with list of Asian spp., P. middlemissi Reed excluded p. 406, Kobayashi (4).

Protobronteus gen. n. [Scutelluidae] for type Eobronteus reeds Sinclair p. 55 (245) Ordovician (Middle) Canada (Quebec), Snajdr Rozpr. Úst. úst. geol. 26 1960.

Protolenid (almost certainly Myopsolenus) p. 103, Basserr in Bassett & Walton.

Protolenidae see Kameschkowiella gen, n.

Protolenidae see Krolina gen. n.

Protolenus atlasensis, P. singularis n. spp. [nom. nuda] p. 81, Hupé (3).

Protolenus planus Pokrovskaya p. 177 pl. i fig. 16, Repina.

Protopliomerops kazachstanicus Balashova [nom. nud.] pl. xi fig. 22, Chernysheva et al.

Protopliomerops punctatus Kobayashi [may belong to Koraipsis] p. 261 pl. xiii figs. 14-15, Kobayashi (3).

Protoptychopyge gen. n. [Asaphinae] p. 125 for type P. prisca (Lamansky) 1905 for Asaphus priscus L. lectotype (Lesnikova 1949 pl. 69 figs. 1a, b) selected and figured including in synonymy Ptychopyge plautini of Schmidt 1907 partim (pl. 3 fig. 15) p. 126 pl. vii figs. 1-5 Ordovician (Lower) U.S.S.R. (Baltio), Balashoya Paleont. Zhurnal 1959 4 1960.

Protoscutellum gen. n. [Scutelluidae] p. 63 (247) for type Bronteus simulans Barrande 1852 [lectotype chosen and figured] excluding B. pl. 48 figs. 31-32 (to Kosvoopeltis partschi p.) p. 64 pl. ii fig. 7 Silurian Bohemia, Snajdr Rozpr. Ust. úst. geol. 26 1960.

Pruvostina referred to Bigotinopsidae fam. n. p. 77, Hupé (3).

Pseudagnostus Jackel 1909 discussed with remarks on pygidial morphology p. 7, P. cavernosus sp. n. p. 12 pl. i figs. 1-4, P. vulgaris sp. n. p. 14 pl. i figs. 5-13 text-fig. 4 Cambrian (Upper) U.S.S.R. (Salair), Rozova Trud. Inst. geol. geophiz. AN S.S.S.R. (Sib. Otd.) 5 1960.

Pseudagnostus Jackel—some Resser 1938 spp. of Oedorhachis [quod vide] may belong p. 13, Poulsen (2).

Pseudagnostus chinensis (Dames) p. 340, P. cyclopygeformis (Sun) excluding Endo 1937 partim p. 341 pl. xix figs. 6-7, P. douvillei (Bergeron) p. 333, Kobayashi (4).

Pa field

(1

P

pl. i

p. 6

Pa varie 1-5, Pa p. 24

Ps (3).

Henr

Ps ascril densi with

(Low

Ps
6a, I
pl. xi
P.
textgen.

Pareferra

Pa Hupé Pa eludir figs.

synor

Ps. S. Schré (for 1

Pariof Policy (Upper

Pse (4).

n

of.

n.

6,

n.

1).

or

b)

0.

5)

R. 0.

7)

an

n.

n.

R.

of en Pseudagnostus communis (Hall & Whitfield) p. 61 pl. iv figs. 3-4, P. prolongus (H. & W.) pygidium newly referred [excluding Palmer 1955 pl. 19 fig. 18 to P. sp.] p. 61 pl. iv figs. 5-6, P. ? acutus (Kobayashi) lectotype selected and figured for Homagnostus a. K. p. 62 pl. iv figs. 10-12, Palmer (1).

Pseudagnostus convergens p. 43, Lochman-Balk.

Pseudagnostus sp. cf. P. prolongus (Hall & Whitfield) p. 812 pl. xev fig. 36, Lochman & Hu.

Pseudonsaphus limatus Jaanusson 1953 pygidial variation discussed p. 245 pl. ix fig. 4, pl. xiii figs. 1-5, P. sp. p. 247 pl. xiii fig. 6, Henningsmoon.

Pseudobasilicus cf. kuckersianus (Schmidt 1898) p. 247 pl. xiii fig. 7. P. cf. kegelensis (S. 1898) p. 247 pl. xiv figs. 1, 4, ? 8, P. ? sp. p. 248 pl. xiv figs. 2-3, Henningsmoen.

Pseudocobboldia n. gen. [nom. nud.] p. 76, Hupé (3).

Pseudoeteraspis angarensis Chernysheva 1950 [not seen] p. 214 pl. v figs. 10-14, pl. viii figs. 4-5 Cambrian (Lower) U.S.S.R. (E. Sayan), Repins.

Pseudogygites latimarginatus (Hall) description of ascribed degree 3 meraspis p. 1118 pls. i-ii, P. canadensis (Chapman) [type of P.] probably synonymous with P. l. p. 1120, Fritz.

Pseudokainella iwayai Kobayashi p. 245 text-fig. 6a, P. a [α] sp. K. 1953 p. 246, P. f sp. nov. p. 246 pl. xii fig. 1 text-fig. 5d [where called P. f sp. indt.], "P." maladiformis (K.) for Mansuyia m. K. p. 242 text-fig. 5e; P. f β sp. K. 1953 referred to Antiaspis gen. n. p. 248, Kobayashi (3).

Pseudokainella? macarenae Harrington & Kay referred tentatively to Poletaevia gen. n. p. 39, Rozova (2).

Pseudokoldinioidia Endo 1944 discussed and referred to Dorypygidae p. 349, Kobayashi (4).

Pseudolenus tenuifrons n. sp. [nom. nud.] p. 81, Hupé (3).

Pseudomegalaspis patagiata (Törnquist 1884) including P. formosa of Jaanusson 1953 partim (pl. 10 figs. 2-3) p. 248 pl. xii figs. 1-7, relations and possible synonymy of P. p. and P. f. p. 249, P. ? sp. p. 251 pl. xii fig. 8, **Henningsmoen**.

Pseudophillipsia croatica Simič sp. n. for Ps. n. sp. S. 1938 p. 31 (33) Carboniferous (Upper) Croatia, Schréter Földt. Közl. 78 1948 [omitted Z. R. 90 11 (for 1952) 1955].

Pseudophillipsia croatica Schréter 1948 a synonym of Paladin mucronatus p. 47n, Bouček & Přibyl.

Pseudosaratogia leptogranulata sp. n. p. 101 pl. xi fig. 10, P. abnormis sp. n. p. 101 pl. xi fig. 11 Cambrian (Upper) U.S.A. (Nevada), Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Pseudosolenopleura Sun discussed p. 378, Kobayashi (4).

Pseudosphaerexochus octacanthus (Angelin 1854) for Cyrtometopus o. A. restr. [non pl. 22 fig. 6] lectotype [fig. 6a] selected and refigured p. 134 pl. xxv fig. 7, P. wolkne sp. n. p. 134 pl. xxiv figs. 5a-b, pl. xxv figs. 1-2, text-fig. 37, P. laticeps (Linnarsson 1869) recorded p. 137, P. sp. a p. 137 pl. xxiv figs. 6-7, P. sp. b p. 138 pl. xxv figs. 6a-c, P. sp. c p. 139 pl. xxii fig. 1 Ordovician (Upper) Poland (Lysa Gora), Kielan Palacont. polon. No. 11 1960.

Pseudotermierella n. gen. [nom. nud.] [Termierellidae] p. 77, Hupé (3),

Pterocephalia Roemer 1849 discussed and variation analysed p. 86, ventral cephalic sutures text-fig. 84, P. concava sp. n, p. 88 pl. ix figs. 1-6, 9-12, P. elongata sp. n, p. 88 pl. ix figs. 14-20 Cambrian (Upper) U.S.A. (Nevada), P. sanctisabae R. with all Resser's 1938 spp. (including P. bridgei) as synonyms p. 88 pl. ix figs. 7-8, 13 text-fig. 19 (reconstr.), P. occidens Walcott holotype figured and considered indeterminate p. 88 pl. ix fig. 21, P. cf. P. e. for P. cf. P. o. of Palmer 1954, Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Pterocephalia bridgei Resser p. 814 pl. xcvi figs. 28-33, Lochman & Hu.

Pterocephaliidae discussed and classified into Aphelaspidinae subfam. n. and Pterocephaliinae; Pterocephalops, Nericia and Grönwallia may belong p. 80, Palmer (1).

Pterocephaliidae see Anechocephalus gen. n.

Pterocephaliidae see Hypostomes A-D.

Pterocephaliidae (Aphelaspidinae) see Taenora gen. n.

Pterocephalinae discussed p. 84 and taken to include Pterocephalia, Cernuolimbus and Sigmocheilus genn. n. and possibly Anechocephalus gen. n., Nericia septemdentata and Dikelocephalites flabelliformis Sun, Palmer (1).

Pterocephaliinae see Cernuolimbus gen. n.

Pterocephaliinae see Sigmocheilus gen, n.

Pterocephalina grata, notha, pogonipensis, utahensis Resser 1942 spp. referred to Sigmocheilus gan. n. with P. cf. P. g. of Wilson 1954, P. tezzna R. probably referable to Labiostria p. 90, Palmer (1).

Pterocephalops may belong to Pterocephaliidae p. 80, Palmer (1).

Pterygometopinae discussed p. 8, Pillet (3).

Pterygometopinae see Liocnemis gen. n.

Pterygometopus Schmidt 1882 discussed p. 8 and classified into P. (Pterygometopus), P. (Achatella), P. (Truncatometopus) subgen. n. p. 9 for type Phacops truncatocaudatus Portlock 1843 text-fig. 4b also including trigonocephalus, kuckersiana, nieszkowskii and kegelensis Schmidt spp.; P. (P.) dagon Reed 1915 [including Dalmanitina proaeva of Bordet et al. 1959 p. 5n] p. 9 pl. i figs. 1-3 text-fig. 5 b, Pillet Bull. Soc. géol. Fr. (7) 2 1 1960.

Ptychagnostus punctuosus (Angelin) p. 10, Noakes et al.

Ptychagnostus sp. text-fig. 4j, Whittington (3).

Ptychocheilus discretus table 1 p. 6, Volk.

Ptychometopinae subfam. n. [Pharostomidae] for Ptychometopus Schmidt p. 161, Balashova Principles of Palaeontology. Arthropods—Trilobitomorphs and Crustaceans 1960.

Ptychometopus referred to Ptychometopinae subfam. n. p. 161, Chernysheva et al.

Ptychometopus—Endocrania compared p. 41, Kobayashi (1).

Ptychoparia striata (Emmrich) text-fig. 536, Müller.

Ptychopariida—list of suborders including Calymenina p. 233, Kobayashi (3).

Ptychopariidae discussed p. 127, Suvorova.

Ptychopariidae—Lermontoviella Pokrovskaya referred p. 201, Repina.

Ptychoparioidea see Anechocephalus gen. n.

Ptychoparioidea see Minupeltis gen, n,

Ptychoparioidea see Morosa gen. n.

Pumilina gen. n. [Aldoniidae] for type P. pandica sp. n. p. 220 pl. xv figs. 13-15 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S. S.S.R. 4 1960.

Quadraticephalus Sun discussed with list of referred spp. p. 401, Q. coreanicus sp. n. for Q. manchuricus of Kobayashi 1935 partim (pl. 6 figs. 1-5) p. 402 pl. xx figs. 2-6, [7], pl. xxi fig. 13, text-figs. 12d-e, Q. quadratus K. p. 402 pl. xix fig. 11, Q. elongatus K. for Q. m. of K. 1935 partim (pl. 6 figs. 6-7) Cambrian (Upper) S. Korea, Q. m. pl. xix fig. 13, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Raaschella Lochman 1938 silently synonymised with Glaphyraspis Resser 1937 p. 279, Rasetti in Harrington et al. [omitted Z.R. 96 11].

Raaschella occidentalis Lochman referred to Glaphyraspis p. 815, Lochman & Hu.

Radiaspis radiata (Goldf.) p. 350, Chlupáč & Vaněk.

Raphiophoridae see Ampyxella gen. n.

Raphiophorus see Rhaphiophorus.

Raphiophorus tenellus (Barrande 1872) p. 165 pl. xxxv fig. 6, R. gratus (B. 1872) with synonyms Ampyx globifrons Olin 1906 [and Dindymene venusta O. refigured p. 147] p. 166 pl. xxxii fig. 6, pl. xxxii figs. 4-5, pl. xxxvi fig. 7, text-fig. 48, R. acus (Troedsson 1918) [may be a synonym of R. setirostris Angelin] p. 168 pl. xxxii figs. 4a-b, pl. xxxv fig. 7, R. cf. a. p. 169; R. depressus A. a synonym of Lonchodomas portlocki (B.) p. 171, Kielan (1).

Raymondella nordica sp. n. p. 37 [pl. vi figs. 15-16] Ordovician (Middle) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Reacalymene limba Shirley p. 158, Whittard.

? Reacalymene pusulosa Shirley p. 167, Dean (3).

Redlichia saitoi Lu p. 343, Kobayashi (4).

Redlichia ? sp. p. 10 pl. i fig. 10, Poulsen (1).

Redlichina vologdini Lermontova p. 177 pl. xiii fig. 5, R. exacuta sp. n. p. 177 pl. xiii figs. 1-4 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Reedocalymeninae discussed and referred to Synhomalonotidae fam. n. p. 45, Kobayashi (1).

Remopleurides text-figs. 3 A-C, Whittington (2).

Remopleurides l' sp. a p. 67 pl. iii fig. 10, R. l' sp. b p. 68 text-fig. 16 Ordovician (Upper) Poland (Lysa Gora), Kielan (1).

Resseropsis gen. n. [nom. nud.] [Neoredlichiidae] p. 76, Hupé (3).

Reuscholithus reuschi Bancroft holotype figured and immature cephala described p. 91 pl. xiv figs. 1-14, Dean (1).

Rhaphiophorus [sic] sp. p. 116, Lawson.

Richardsonella flabellifera (Hall & Whitfield) discussed p. 243, Kobayashi (3).

Richardsonellidae Raymond discussed p. 239, Apatokephalina probably referable p. 242, Kobayashi (3).

Rincónia pardailhensis, R. angusta n. spp. [nom. nuda] p. 82, Hupé (3).

Rochaeva n. gen. [nom. nud.] [Termierellidae] p. 77, Hupé (3).

Rondocephalus mirandus Pokrovskaya p. 205 pl. vii figs. 8-9, Repina.

Rudagnostus referred to Geragnostidae p. 234, Kobayashi (3).

Sajanaspis gen. n. [Dolerolenidae] p. 196 for type S. pokrovskajae sp. n. p. 197 pl. xi figs. 1-2, S. modesta sp. n. p. 197 pl. xi fig. 4, S. sp. pl. xi fig. 3, S. sp. pl. xii fig. 8 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Salterolithus Bancroft 1929 discussed p. 95, S. caractaci (Murchison) including S. sp. nov. Dean 1958 p. 96 pl. xv figs. 1-2, 4, 6, 9, S. praecursor 5p. n. including S. sp. B. 1929 p. 98 pl. xv figs. 3, 5, 7-8, 10-12 Ordovician (Middle) Britain (Shropshire); S. smeathenensis B. 1949 type of Smeathenia gen. n. p. 100, S. harnagensis B. referred to Broeggerolithus p. 105, Dean Bull. Brit. Mus. (nat. Hist.) [Gool.] 4 4 1960.

Sao hirsuta Barrande text-fig. 547, Müller.

Koba Koba Sar p. 40

[19

Sat

San

Suvo

Sea

140 (2

chose 1-3, 1 and lector pl. x: subsp 8. hose elonge figs. turdis xxvii p. 15 furcif

with

holot

figs. Bohe

Maks
Sch

n. p.

Repi

Sou 1925 Snaje

Sen Metascutei

Sci S. et figd. S. la Kolii Spin P. of S. b

8. b inern pusts ).

an

to

2).

nd

e]

d)

m.

7,

ol.

4,

ì),

m

);

Saukia Walcott discussed with list of Asian spp., S. vagans Reed excluded p. 405; S. sp. listed p. 335, Kobayashi (4).

Saukiella sp. for Calvinella sp. of Kobayashi 1935 p. 407 pl. xix fig. 30, Kobayashi (4).

Saukiinae discussed p. 403, Kobayashi (4).

Saukiinae see Platysaukia gen. n.

Sayanella Pokrovskaya 1959 a synonym of Botomella Suvorova 1958 p. 149, Suvorova.

Scabriscutellum see Scutellum.

Scabriscutellum R. & E. Richter 1956 discussed p. 140 (256), S. oblongum (Hawle & Corda 1847) lectotype chosen and figd. p. 142 pl. xxv figs. 12-14, pl. xxv figs. 1-3, 10, S. caelebs c. (Barrande 1852) lectotype chosen and figd. with synonym Bronteus gigas H. & C. lectotype chosen and figd. p. 143 pl. xxvi figs. 4-9, pl. xxxvi fig. 13, text-fig. 46, S. caelebs consimile subsp. n. p. 145 (256) pl. xxvi fig. 11, pl. xxvii fig. 15, S. holynense sp. n. p. 146 (257) pl. xxvi fig. 12, S. elongatum (B. 1846) holotype figd. p. 147 pl. xxvii figs. 11-12, S. richteri (B. 1852) with synonym B. xxvii figs. 12-3, S. billingsi (B. 1872) holotype figd. p. 148 pl. xxvii figs. 2-3, S. billingsi (B. 1872) holotype figd. p. 150 pl. xxviii fig. 13 text-fig. 4, S. (nov. subg. ?) furciferum (H. & C. 1847) lectotype chosen and figd. with synonym Scutellum analogum Prantl 1949 holotype figd. p. 152 pl. xxviii figs. 4-12, pl. xxviii figs. 1-7 text-fig. 47, S. sp. pl. v fig. 1 Devonian (Middle) Bohemia, Snajdr Rozpr. Ust. üst. geol. 26 1960.

Scabriscutellum sibiricum (Tschernyschew 1893) holotype figured p. 32 pl. i figs. 2-4, pl. iv fig. 4, S. aff. scabrum Goldfuss 1843 p. 34 pl. i fig. 5, Maksimova.

Schistocephalus sibiricus sp. n. p. 214 pl. xvi figs. 10-11 Cambrian (Middle) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Scinocephalus referred to Proapatokephalidae fam. n. p. 33, Rozova (2).

Scotiella sp. p. 350, Tomczykowa,

Scutelluidae (Hawle & Corda 1847) R. & E. Richter 1925 discussed and *Dulanaspis* excluded p. 53 (245), Snajdr.

Scutelluidae R. & E. Richter preferred [to Thysanopeltidae] p. 415, Whittington (1).

Scutelluidae see Breviscutellum, Cornuscutellum, Metascutellum, Microscutellum, Protobronteus, Protoscutellum, Spiniscutellum genn. n.

Scutellum Pusch 1833 discussed p. 181 (259), S. edwardsi (Barrande 1852) lectotype chosen and fgd. p. 182 pl. vii fig. 10, pl. ix figs. 6-12, text-fig. 56; S. lanceolatum Prantl & Přibyl a synonym of Kolihapeltis parabolina p. 165, S. přibyli Prantl of Spiniscutellum umbelliferum p. 113, S. analogum P. of Scabriscutellum furciferum p. 152, S. nováki & S. bohemicum Prantl & Přibyl of Breviscutellum inerme p. 174, S. gaudryi P. & P. of Metascutellum pustulatum p. 185, Snajdr.

Scutellum alutaceum Goldfuss 1843 p. 27 pl. viii fig. 7, S. longicaudatum Maksimova 1955 p. 29 pl. viii figs. 1-5, S. elegans Peetz [Petts] 1901 p. 30 pl. iii fig. 9, S. cf. orientalis M. 1955 p. 31 pl. viii fig. 6, Maksimova.

Scutellum (S.) brevifrons (Barr.), S. (S.) oblongum (H. & C.), S. (Scabriscutellum) elongatum (B.) p. 350, Chlupáč & Vaněk.

"Soutellum" multiverrucatum sp. n. for Bronteus pustulatus of Barrande 1852 partim (pl. 48 figs. 13-16) and 1872 p. 202 (262) pl. xxxv figs. 1-5, "S." obscurum sp. n. p. 203 (262) pl. xxxv figs. 6-7, "S." asperulum (B. 1872) holotype figd. p. 204 pl. xxxv figs. 8-10, "S." setosum Frantl 1949 holotype figd. p. 205 pl. xxxv figs. 17, "S." schöbli (Novák 1884) lectotype chosen and figd. p. 206 pl. xxxv figs. 11-13, "S." sieberi (Hawle & Corda 1847) lectotype chosen and figd. p. 208 pl. xxxvi figs. 1-2 Devonian (Lower) Bohomia, **Snajdr** Rozpr. Ust. úst. geol. **26** 1960.

Scutellum paliferum (Beyrich) text-fig. 564 B, Müller.

Scutellum (Thysanopeltis) speciosum speciosum (Hawle et Corda 1847) em. Prantl 1950 p. 336 pl. ii fig. 11, Strnad (1).

Scutellum viator Barr. p. 45, Cogney.

Scutellum sp. p. 160, Cavet.

Scutellum sp. indet. p. 269, Churkin & Langenheim.

Seleneceme Clark 1924 discussed p. 117, S. acuticaudata (Hicks) for Illaenopsis ? a. H. holotype figured p. 118 pl. xvi figs. 1-5, Whittard.

Semiproetus 800 Proetus.

Serksia n. gen. [nom. nud.] [Olenellid] p. 76, S. cordiformis n. sp. [nom. nud.] p. 81, **Hupé** (3).

Serksioides n. gen. [nom. nud.] [Olenellid] p. 76, Hupé (3).

Serrodiscus see Eodiscid?

Shantungia (Metashantungia) Chang 1959 [not seen] discussed p. 353, S. (M.) brevica C. text-fig. 2q. S. (Parashantungia) Chu 1959 [not seen] discussed p. 353, S. (P.) elongata C. text-fig. 2 p, Kobayashi (4).

Shirakiella Kobayashi 1935 discussed with synonym Mansuyella Endo p. 390, S. ? sp. p. 390 pl. xxi [sio] fig. 12, S. elongata K. pl. xix fig. 26, S. laticonvexa K. p. 334, Kobayashi (4).

Shivelicus parvus Pokrovskaya p. 175 pl. vii fig. 11, Repina.

Shumardia pellizzarii Kobayashi 1934 p. 235 pl. xiii figs. 23-25, Kobayashi (3).

Shumardia polonica sp. n. p. 159 pl. xxxii fig. 5 text-fig. 45 Ordovician (Upper) Poland (Lysa Gora), Kielan Palaeont. polon. No. 11 1960.

Shumardia pusiliformis Poletaeva [not seen] pl. iv fig. 20, Chernysheva et al.

Sigmocheilus gen. n. [Pterocephaliinae] p. 89 for type S. serratus sp. n. p. 91 pl. x figs. 1-3 text-fig. 20 (reconstr.), S. grata (Resser) for Pterocephalina g. R. p. 90 pl. ix figs. 22-23, 26-27, S. pogonipensis R. for P. p. R. p. 91 pl. x figs. 4-7, S. utahensis (R). for P. u. R. p. 91 pl. ix figs. 24-25, 28 Cambrian (Upper) U.S.A. (Nevada); P. notha R., P. of. P. g. of Wilson 1954 and Iddingsia ? quinnensis R. also referred to S. p. 90, Labiostria sigmoidalis Palmer related to S. p. 80, Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Sinosaukia Sun discussed p. 404, Kobayashi (4).

Sinskia gen, n. [Holmiinae] p. 18 for type S. optabilis sp. n. p. 19 pl. i figs. 1-4 text-fig. 2 Cambrian (Lower) U.S.S.R. (Siberian Platform), Suvorova Trud. paleont. Inst. S.S.S.R. 34 1960.

Smeathenia gen. n. [Trinucleidae] for type Salterolithus smeathenensis Bancroft 1949 lectotype selected and figured p. 100 pl. xiii figs. 7, 10-16 Ordovician (Middle) Britain (Shropshire), Dean Bull. Brit. Mus. (nat. Hist.) [Geol.] 4 4 1960.

Solenoparia Kobayashi with discussion of referred spp. p. 379, S. laevis sp. n. p. 380 pl. xix figs. 14 a-b, 15-16 Cambrian (Middle) S. Korea, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Solenopariinae discussed p. 378, Kobayashi (4).

Solenopleurella kolbinica sp. n. p. 200 pl. v figs. 7-9 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Solenopleurellinae see Litaspis gen. n.

Solenopleuridae discussed p. 376, see also Menocephalitinae subfam. n. p. 380, Kobayashi (4).

Spathacolymene gen. n. [Calymeninae] p. 892 for type Calymene nasuta Ulrich 1879 lectotype selected and figured p. 893 pl. cxvi figs. 1-12 Silurian (Middle) U.S.A. (Indiana), Tillman J. Paleont. 34 5 1960.

Sphaeragnostus gaspensis europeensis subsp. n. p. 56 pl. i figs. 1-3 text-fig. 11 Ordovician (Upper) Poland (Lysa Gora), Kielan Palaeont. polon. No. 11 1960.

Sphaerexochus aff. mirus Beyrich 1845 p. 11 pl. i figs. 8-9, text-fig. 5e, Pillet (3).

Sphaerexochus taimyricus sp. n. p. 29 [pl. v figs. 8 a-6] Ordovician (Upper) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 17 [pls. in 14] 1960.

Sphaerexochus sp. p. 31, Bogdanov & Chugaeva.

Sphaerocoryphe Angelin 1854—proposed designation of S. dentata A. 1854 as type p. 232, Stubblefield & Henningsmoen.

Sphaerocoryphe ef. thompsoni (Reed) p. 334, Cave.

Sphaerometopus Angelin 1854—proposed suppression p. 232, Stubblefield & Henningsmoon.

Spiniscutellum gen. n. [Scutelluidae] p. 111 (254) for type S. umbelliferum (Beyrich 1845) for Brontess umbellifer B. neotype chosen and figd. [excluding Prantl & Přibyl 1947 partim (pl. 6 fig. 2 to Decoroscutellum hanuší) and with synonyms B. sparens Hawle & Corda (lectotype chosen and figd.) and Scutellum přibyli Prantl (holotype figd.) and including S. partschi of Howell 1961] p. 112 pl. viii figs. 1-19, pl. ix figs. 1-4, pl. xxxvi fig. 12, text-figs. 5, 39-40 Silurian (Upper) and Devonian (Lower) Bohemia, Snajdr Rozpr. Úst. úst. geol. 26 1960.

Staurocephalus clavifrons Angelin 1854 p. 156 pl. xxiv fig. 9, pl. xxvi figs. 7-9, Kielan (1).

Stigmagnostus gen, n. [Agnostidea (? Peronopsinae)] for type ? Tomagnostus canotensis Rusconi 1951 [re-interpreted as a pygidium] p. 15 pl. i figs. 11-12 Cambrian (Middle) Argentina (Mendoza), Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Strenuella strenua (Billings) from Morocco p. 77, Hupé (3).

Stubblefieldia sp. p. 139 pl. xiii figs. 5a-b, pl. xxiv fig. 8, pl. xxv figs. 8a-b; S. neglecta of Barrande 1872 partim [pl. 3 fig. 18] probably belongs to Actinopeltis insocialis (B.) p. 140, Kielan (1).

Stygina sp. p. 30, Bogdanov & Chugaeva.

Stygina sp. (conspecific with Skjeseth 1955 pl. 2 fig. 3) p. 82 pl. viii figs. 1-2, Kielan (1).

Symphysops subarmata elongata subsp. n. p. 90 pl. xi figs. 1-4, S. sp. a [may belong to S. s. e.] p. 92 pl. xi fig. 6, S. sp. b p. 92 pl. xi fig. 5 Ordovician (Upper) Poland (Lysa Gora), Kielan Palaeont. polon. No. 11 1960.

Symphysurina brevispicata p. 43, Lochman-Balk.

Symphysurus angustatus Sars & Boeck pp. 216, 217, Tjernvik.

Symphysurus convexus Balashova [nom. nud.] pl. vi figs. 20-21, Chernysheva et al.

Symphysurus superstes Olin 1906 restr. lectotype selected and a synonym of Microparia speciosa [q.v.], S. s. O. partim (pl. 3 fig. 6) referred to Panderia megalophthalma p. 90, S. s. of Wiman 1907 is an Illaenid p. 90, Kielan (1).

Synhomalonotidae fam. n. [Calymenacea] p. 42 for Synhomalonotinae and Reedocalymeninae, Kobayashi Japan. J. Geol. Geogr. 31 1 1960.

Synhomalonotinae subfam. n. [Synhomalonotidae] for Synhomalonotius and Victnamia gen. n. p. 42, Kobayashi Japan. J. Geol. Geogr. 31 1 1960.

Synhomalonotinae see Vietnamia gen. n.

Synhomalonotus Pompeckj 1898 a synonym of Neseuretus Hicks [q.v.] p. 139, Whittard.

Synhomalonotus Pompeckj discussed with list of spp. and referred to Synhomalonotidae (Synhomalonotinae) subfam. n. p. 42, "S." birmanicus (Reed) text-fig. 2e; S. kobayashii Harrington & Leanza may not belong p. 45, Kobayashi (1).

Syn. 1538,

[196

Majew Tae Elburg

Tae stages Shaw T. sp.

p. 83 20-23 U.S.A tative geol.

Tai group Tai

maner
pl. x
Manel
Univ.

T. wo

Take type fig. 2, sp. n Arger dansk

Ta: p. 77, Tel p. 46 Koba;

Coron (Low 8.S.S

Te

Th 156 ( and Barra redier pl. xi text-i

p. 16

g

d

12

7,

is

2

an

n.

k.

7,

ol.

pe

ia

an

42

of

(be

Synhomalonotus fedalensis n. sp. [nom. nud.] p. 1538. Destombes et al.

Symphoroides sp. pl. vi figs. 15-17, Wilson & Majewske.

Taenicephalus quinnensis Resser referred to Elburgia gen. n. [q.v.] p. 69, Palmer (1).

Taenicephalus shumardi (Hall) with immature stages and including T. cordillerensis of DeLand & Shaw 1956 (pl. 67 fig. 3) p. 811 pl. xev figs. 12-23, 31, T. sp. undet. p. 798, Lochman & Hu.

Taenora gen. n. [Pterocephaliidae (Aphelaspidinae)] p. 83 for type T. expansa sp. n. p. 84 pl. vii figs. 20-23 text-fig. 17 (reconstr.) Cambrian (Upper) U.S.A. (Nevada); Labiostria platifrons Palmer tentatively referred to T. p. 84, Palmer Prof. Pap. U.S. geol. Surv. No. 334-C 1960.

Taihungshania Sun discussed and Chinese spp. grouped into two series p. 256, Kobayashi (3).

Taihungshaniidae discussed p. 256, Kobayashi (3).

Taipaikia gen. n. [Tsinaniidae] for type Paramansuyella glabra Endo p. 398 pl. xxi figs. 20-21, ? pl. xx fig. 18, text-fig. 11h Cambrian (Upper) Manchuria and Korea, Kobayashi J. Fac. Sei. Tokyo Univ. [2] 12 2 1960.

Taitzehoia Chu 1959 [not seen] discussed p. 352, T. wangi C. text-fig. 2g, Kobayashi (4).

Taitzuia Resser & Endo referred to Ordosiinae and spp. discussed p. 371, Kobayashi (4).

Talbotina see Genus and species undet. 2.

Talbotinella gen. n. [Lonchocephalidae] p. 23 for type T. communis sp. n. p. 24 pl. ii figs. 2-8, pl. iii fig. 2, T. leanzai sp. n. p. 25 pl. ii figs. 9-12, T. rusconsi sp. n. p. 27 pl. ii figs. 13-15 Cambrian (Middle) Argentina (Mendoza), Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Tazemmourtia n. gen. [nom. nud.] [Bigotinopsidae] p. 77, Hupé (3).

Tellerina Ulrich & Resser with list of Asian spp. p. 406, T. coreanica Kobayashi listed p. 334, Kobayashi (4).

Terechtaspis gen. n. [Komaspidae] for type T. coronaria sp. n. p. 208 pl. xiii figs. 6-7 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.R. 4 1960.

Terranovella see Genus and species undet. 1.

Thaleops sp. p. 29, Bogdanov & Chugaeva.

Thysanopeltis Hawle & Corda 1847 discussed p. 186 (257), T. speciosa H. & C. 1847 neotype chosen and figd. [with synonyms Bronteus thysanopeltis Barrande 1852, Scutellum (T.) s. abreviatum and s. redivirum Prantl 1949 both holotypes figd.] p. 157 pl. xxix figs. 4-8, pl. xxx figs. 1-14, pl. xxxi figs. 1-8, T. clementina (B. 1872) holotype figd. p. 162 pl. xxix figs. 1-2 text-fig. 49, Snajdr.

Thysanopeltis paucispinosa sp. n. p. 137 pl. i figs. 5a-c Devonian (Middle) Japan (Iwate), Okubo Earth Science [Chikyū Kagaku] No. 4 1951.

Thysanopeltis see Scutellum.

Tingocephalinae Hupé referred provisionally to Kaolishaniidae p. 355, Kobayashi (4).

Tingocephalinae see Kabutocrania gen. n.

Tingocephalus Sun discussed, T. concavolimbatus (Endo) discussed p. 357, T. magnus sp. n. p. 358 pl. xx fig. 17 Cambrian (Upper) S. Korea, Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Tolstotchichaspis gen. n. [Plethopeltidae] p. 59 for type T. grandis sp. n. p. 60 pl. vii figs. 11-20 Cambrian (Upper) U.S.S.R. (Salair), Rozova Trud. Inst. geol. geophiz. AN S.S.S.R. (Sib. Otd.) 5 1960.

? Tomagnostus canotensis Rusconi reinterpreted as pygidium and type of Stigmagnostus gen. n. [q.v.] p. 15, Poulsen (2).

Tonkinella orientalis Kobayashi p. 212 pl. xvii figs. 9-10, T. sisimica sp. n. p. 212 pl. xvii figs. 1-8 Cambrian (Middle) U.S.S.R. (E. Sayan), Repina Regional Strat. S.S.R. 4 1960.

Tretaspis granulata (Wahlenberg 1818) restr. type refigured with synonyms Trinucleus bucklandi Barrande restr. and T. elliptifrons Olin p. 171 pl. xxxiii figs. 1-3, pl. xxxiv figs. 1-2, pl. xxxvi fig. 6, text-fig. 49 (graph), T. seticornis seticornis Stermer 1945 [sie] including T. b. B. 1852 partim (pl. 29 fig. 10, pl. 30 figs. 14-16) p. 174 pl. xxxiii fig. 6, Kielan (1).

Tretaspis moeldenensis sp. n. p. 334 pl. x figs. 1-7 Ordovician (Upper) Britain (S. Wales); T. kiaeri of King & Williams 1948 [non fig.] and T. aff. k. of Lamont 1945 are similar to T. m., T. of. k. of Whittington 1938 may be new sp. p. 336, Cave Geol. Mag. 97 4 1960.

Trierepicephalidae see Prometeoraspis gen. n.

Tricrepicephalus sp. cf. T. coria Walcott p. 817 pl. xevii figs. 21-26, T. sp. undet. p. 816, Lochman & Hu.

Trilobites asellus Esmark referred to Ogmocnemis gen. n. p. 69, Kielan (1).

Trilobites sp. [evidently new genus, probably new family] p. 178 pl. iii figs. 4-5, pl. xxi fig. 3 Ordovician (Upper) Poland (Lysa Gora), Kielan (1).

Triniidae referred to Corynexochoidea p. 78, Chernysheva et al.

Trinodus tardus (Barrande 1846) with synonym Agnostus glabratus Angelin p. 59 pl. i figs. 6-13 text-fig. 13, T. sp. p. 62 pl. i fig. 5 text-fig. 14; Dindymene minuta Reed 1931 may be part of a Trinodus p. 147, Kielan (1).

Trinucleid p. 141, Cavet.

Trinucleidae discussed with list of genera p. 84, evolution in Shropshire p. 138, Dean (1).

Trinucleidae see Smeathenia gen, n.

Trinucleus bucklandi Barrande and T. elliptifrons Olin synonyms of Tretaspis granulata (Wahlenberg) p. 171, Kielan (1).

Trinucleus discors Angelin may belong to Broeggerolithus p. 104, T. gibbifrons McCoy suggested that name be rejected, T. g. of Reed 1914 resembles B. globiceps p. 104 [see also Cryptolithus], Dean (1).

Trinucleus sp. p. 31, Bogdanov & Chugaeva.

Trochurus sp. indet. p. 269, Churkin & Langenheim.

" Trochurus" sp. p. 350, Tomczykowa.

Tropidocoryphe filicostata (Novák 1890) p. 337 pl. iii figs. 12-13, Strnad (1).

Tropidocoryphinae—Phyllaspis tentatively referred p. 119, Alberti.

Tropidopyge Harrington & Kay discussed with implied reference to Birmanitidae fam. n. p. 256, Kobayashi (3).

Truncatometopus subgen. n. see Pterygometopus.

Tsinania canens (Walcott) p. 398, T. ceres (W.) p. 335, Kobayashi (4).

Tsinaniidae discussed and Esseigania, Koldiniella, Kazelia, Jubileia may belong p. 396, Kobayashi (4).

Tsinaniidae see Taipaikia gen. n.

Tumulina gen. n. [Dinesidae] p. 204 for type T. tumidica sp. n. p. 205 pl. xii figs. 1-3 Cambrian (Lower) U.S.B. (E. Sayan), Repina Regional Strat. S.S.S.R. 4 1960.

Tunqueella gen. n. [Neoredlichiidae] p. 178 for type T. manica sp. n. p. 179 pl. iv figs. 5-9, T. obesa sp. n. p. 181 pl. iv figs. 10-14 Cambrian (Lower) U.S.S.R. (E. Sayan), T. nitens and T. procera (Suvorova spp.) transferred from Bergeroniaspis p. 179, Repina Regional Strat. S.S.S.R. 4 1960.

Ullaspis [Westergård] may belong to Housiidae p. 74, Palmer (1).

Ulricholithus Bancroft 1949 a synonym of Broeggerolithus pp. 86, 102, Dean (1).

Uncaspis sp. undet. p. 816 pl. xevii fig. 14, Lochman & Hu.

Unguliproetus see Proetus.

Vietnamia gen. n. [Synhomalonotinae] p. 43 for type Calymene dowillei Mansuy 1908 p. 44 pl. v figs. 4.5 text-fig. 2d Ordovician Viet Nam, Kobayashi Japan. J. Geol. Geogr. 31 1 1960.

Warburgaspis Přibyl 1946 considered nom. nud. p. 69, Kielan (1).

Waribole see Cyrtosymbole.

Weberides—range of mucronate and non-mucronate spep p. 122, W. mucronatus (M'Coy), W. m. traquairi Reed, W. parilis cuspidata R. W. aff. shunnerensis (King), W. sp. listed p. 129, Johnson.

Weberides mucronatus (M'Coy) pp. 293, 311 [horizon discussed by W. S. Bisat p. 315; trilobites allied to W. m. recorded by G. A. L. Johnson p. 316], W. barkei (Woodward) p. 311, W. sp. (non-mucronate) pp. 310, 312, Wilson.

Weberopeltis Maksimova 1959 discussed p. 35, W. bublitchenkoi sp. n. p. 36 pl. vi figs. 11-12 text-fig. 1 (4), W. kurjensis sp. n. p. 37 pl. vi figs. 6-8 text-fig. 1 (5) Devonian (Middle) U.S.S.R. (Altai), W. eugeni (Chornysheva) text-fig. 1 (1), W. arcticum (Weber) text-fig. 1 (2), W. yakoolevi (W). text-fig. 1 (3) all so referred pp. 36-37, Maksimova Devonian and Carboniferous trilobites of the Rudny Altai 1960.

Weeksing see Genus and species undet. 2.

Whittingtonia whittingtoni sp. n. p. 109 pl. xvi fig. 5, pl. xviii figs. 1-4, text-fig. 28, W. sp. p. 112 pl. xviii fig. 5 text-fig. 29 Ordovician (Upper) Poland (Lysa Gora), Kielan Palaeont. polon. No. 11 1960.

Williameina gen. n. [Asaphiscinae] p. 18 for type W. cortesi sp. n. p. 19 pl. i figs. 19-24, W. harringtoni sp. n. p. 20 pl. i figs. 29, W. mikkelseni sp. n. p. 21 pl. ii figs. 25-28, W. ornata sp. n. p. 22 pl. ii fig. 1, pl. iii fig. 1 Cambrian (Middle) Argentina (Mendoza), Poulsen Mat.-fys. Medd. danske Vid. Selsk. 32 11 1960.

Yabeia Resser & Endo probably a synonym of Metagraulos p. 387, Kobayashi (4).

Yokusenia Kobayashi 1935—synonymy with Lioparia Lorenz 1906 reaffirmed p. 393, Kobayashi (4),

Yokuseninae replaced by Liopariinae subfam, n. p. 391, Kobayashi (4).

Yosimuraspis gen. n. [Monkaspidae] for type Y. vulgaris 59. n. p. 238 pl. xii figs. 9-20 text-fig. 2 (reconstr.), Y. v. longulum subsp. n. p. 239 pl. xii figs. 7-8 Ordovician (Lower) Korea (South), Kobayashi J. Fac. Sci. Tokyo Univ. [2] 12 2 1960.

Yapingia discussed p. 257, Kobayashi (3).

Zbirovia longifrons (Olin 1906) restr. [? non pl. 3 fig. 16] holotype [sie] refigured p. 97 pl. xii figs. 1-5, Kielan (1).

Zdicella bornholmiensis sp. n. p. 96 pl. xii figs. 7-8, pl. xiii figs. 1-4 text-fig. 24 Ordovician (Upper) Bornholm and Poland, Kielan Palaeont. polon. No. 11 1960.

Zigzuella n. gen. [nom. nud.] [Strenuellinae] p. 77, Hupé (3).

Gen. and sp. indet. A., Gen. and sp. indet. B., Gen. and sp. indet. C (all undescr. pygidia) p. 451 Ordovician (Lower) Spitsbergen, Gobbett & Wilson.

Gen. et sp. indeterminata pl. i figs. 11-12, Gen. et sp. indet. pl. vii fig. 12, Gen. et sp. indet. pl. xii figs. 9-10 p. 223 Cambrian (Lower) U.S.S.R. (E. Sayan), Repina.

Genus and species undet. 1 [compared with Brassicicephalus and Terranovella] p. 825 pl. xovii figs. 38-39, Genus and species undet. 2 [compared with Talbotina and Weeksina] p. 826 pl. xovii fig. 37, Genus and species undet. 3 [compared with Kingstonia and Bynumia] p. 826 pl. xovii figs. 35-36 Cambrian (Upper) U.S.A. (Wyoming), Lochman & Hu.

Palme Genus et sp. pl. ii fi

Argen

[19

Ger

p. 102

Hawle B — I (C.) co Hypoc chosen P. sp. archia

fig. 3,

],

٧.

r)

0.

ni

ii ), 0.

1.

2

Genus and species undetermined 1 p. 101 pl. xi figs. 2, ? 5, 6, Genus and species undetermined 2 p. 102 pl. xi fig. 1, Genus and species undetermined 3 [pygidium, compared with Cliffia latagenae] p. 102 pl. xi fig. 3 Cambrian (Upper) U.S.A. (Nevada), Palmer (1).

Genus et sp. in d. (librigena) p. 32 pl. i fig. 18, Genus et sp. ind. (pygidium) p. 32 pl. ii fig. 26, Genus et sp. ind. (pygidium compared with *Coosella*) p. 33 pl. ii figs. 18-19, Genus et sp. ind. (librigena compared with *Coosia*) p. 33 pl. ii fig. 20 Cambrian (Middle) Argentina (Mendoza), **Poulsen (2)**.

Hypostom A—? Proctus (Cornuproctus) dufrésnoyi Hawle & Corda 1847 p. 213 pl. iv figs. 4-5, Hypostom B—P. (C.) sp. p. 214 pl. iv fig. 6, Hypostom C—? P. (C.) complanatus Barrande 1846 p. 215 pl. iv figs. 1-2, Hypostom D—P. (P.) neglectus B. 1852 [lectotype chosen p. 193] p. 215 pl. iv figs. 7-8, Hypostom E—P. sp. p. 216 pl. iv fig. 9, Hypostom F—Prionopeltis archiact (B. 1846) p. 217 pl. iv fig. 10, Přibyl.

Hypostoma fam. gen. et sp. indt. p. 263 pl. xiv fig. 3, Kobayashi (3).

Hypostome A pl. xi fig. 12, Hypostome B pl. xi fig. 7, Hypostome C pl. xi fig. 4, Hypostome D pl. xi fig. 9, Hypostome E pl. xi fig. 8 p. 60 Cambrian (Upper) U.S.A. (Nevada) [hypostome E assigned to Elviniidae, others associated with Elviniidae and Pterocephaliidae], Palmer (1).

Hypostomes see Proetid hypostomes.

N. gen. aff. Aphelaspis of Palmer & Hazzard 1956 belongs to Prehousia gen. n. p. 78, Palmer (1).

Trilobita indet. table 1 p. 6, Volk.

Trilobite pygidium [probably Phacopoid] p. 40 pl. ix figs. 1-3, Boucot et al.

Trilobites indet. (non-mucronate pygidia) p. 129, Johnson.

Unidentified trilobite fragments p. 103, Bassett & Walton.



